

# ULTRAPULSE CO<sub>2</sub> LASER THERAPY FOR NEVUS SEBACEUS JADASSOHN

A. L. Krunić, R. E. Clark and M. M. Urošević

## ABSTRACT

*Background:* Ultrapulse CO<sub>2</sub> laser could be an effective modality for the ablation of the nevus sebaceus of Jadassohn for cosmetically important areas in selected cases.

*Objective:* Evaluation of effectiveness and cosmetic results concerning the ultrapulse CO<sub>2</sub> laser ablation in 15-year old white boy with nevus sebaceus of the glabellar area.

*Methods:* Ablation of the nevus under local anesthesia with ultrapulse CO<sub>2</sub> laser, in 4 sessions spaced two months apart.

*Results:* An 18-month follow-up disclosed no recurrent diseases with satisfactory cosmetic result.

*Conclusion:* This method produces superior results with minimal risk of scarring compared to a continuous wave CO<sub>2</sub> laser and classical excisional surgery. Although the risk for malignant change of nevus sebaceus on the face is very low, careful follow-ups are mandatory in order to obtain early recognition of any possible malignant change.

## KEY WORDS

*nevus sebaceus of Jadassohn, ultrapulse CO<sub>2</sub> laser ablation, cosmetic results*

---

## INTRODUCTION

Nevus sebaceus of Jadassohn (NSJ) belongs to the group of congenital organoid nevi that characteristically occurs on the scalp and face<sup>1,2</sup>. The overall malignant potential is reported to be between 8-12 %, with the basal cell carcinoma being the most common malignancy developing in these lesions after puberty<sup>3</sup>. The complete surgical excision is considered the treatment of choice, especially for the lesions located

on the scalp which have the highest risk for malignant transformation<sup>4</sup>. The ultrapulse CO<sub>2</sub> laser with high energy and short pulses, achieves controlled penetration, minimal nonspecific thermal damage and char-free ablation<sup>5</sup> and can be a promising tool for the ablation of nevus sebaceus at the cosmetically important locations where conventional surgery may produce significant mutilation and disfigurement.



Figure 1. Nevus sebaceus on the mid forehead prior to treatment with Ultrapulse CO<sub>2</sub> laser.



Figure 2. The treated area immediately after vaporization was completed (4th session).

## CASE REPORT

We present a 15-year old white boy with the 35x15 mm linear nevus sebaceus of the glabellar area. In the infancy the lesion was flatter and pinkish. Recently it became elevated and more yellowish in color (Figure 1). The patient had no history of seizures, eye abnormalities, or impaired intelligence and was in good physical health.

Histopathologic analysis disclosed slightly acanthotic, papillomatous epidermis with increased number of sebaceous gland in the upper half of dermis. The glands were connected to rudimentary follicular infundibula, which were dilated and filled with keratin plugs. These findings were consistent with the diagnosis of nevus sebaceus.

The nevus was ablated under local anesthesia (1% lidocaine with epinephrine) with ultrapulse CO<sub>2</sub> laser (Coherent, Palo Alto, California), in 4 sessions spaced two months apart. The laser settings were 300 mJ energy/pulse, 50W average power, and computerized pattern generator adjusted to the density of 7, sizes from 3 to 5 and patterns of 1 and 3. Each ablation session consisted of 3 passes to the uppermost reticular dermis (Figure 2).

At twelve months after the surgery satisfactory cosmetic result with moderate postinflammatory hyperpigmentation was achieved (Figure 3). The discoloration resolved with combined use of topical hydroquinon 3% solution, tretinoin 0,05% cream and sunscreen. At eighteen-month follow-up no recurrence of the nevus was seen.



Figure 3. One year after ablation. Postinflammatory hyperpigmentation and mild pinkish discoloration is still present.



## DISCUSSION

Nevus sebaceus of Jadassohn (NSJ) is considered to be a localized malformation of epidermis, pilosebaceous units and ectopic apocrine glands which develop in about two thirds of cases<sup>1</sup>. The natural course of NSJ evolves through different stages over time from a deficient malformation (infantile phase) to a hyperplastic sebaceous lesion as a consequence of hormones and sexual maturation (adolescent phase)<sup>1,6</sup>. A third stage commonly develops in adulthood where different benign adnexal tumors arise in NSJ<sup>3</sup>. The most frequent one is believed to be the syringocystadenoma papilliferum, which develops in 8-19% of cases<sup>1,3</sup>. Malignant transformation of NSJ usually occurs in scalp lesions and is considered to be much less frequent. Basal cell carcinoma is the most common malignancy observed in 5-7% of cases<sup>3,4</sup>. In rare instances, squamous cell carcinoma, apocrine carcinoma, and malignant eccrine poroma have developed in NSJ that led to regional and even generalized metastases<sup>4,7</sup>.

Aside from malignant potential, NSJ may present cosmetic or functional problems by virtue of its different locations on the face<sup>2</sup>. For all these reasons, acceptable treatment is often desirable but not always possible. Most investigators consider surgical excision as the treatment of choice<sup>8</sup>. In our patient the tight skin on the forehead altogether with the size of the lesion, would necessitate repair with the advancement flap after the excision. This surgical approach would be cosmetically less satisfactory, since it carries the risk for disfiguring scar with significant impact on

the psychology and self-image of the young adolescent.

Continuous CO<sub>2</sub> laser in nonfocused mode was also used for the ablation of NSJ<sup>9</sup> and epidermal nevi<sup>10</sup>. The ablation of NSJ, although clinically successful, resulted in hypertrophic scar on the nose<sup>9</sup>. Ultrapulse CO<sub>2</sub> laser technology<sup>5</sup> with controlled penetration; char-free ablation and minimal collateral thermal damage produced satisfactory result in our patient, with transitory postinflammatory hyperpigmentation. The superficial nature of the lesion, the lack of the extension to the deep reticular dermis as well as the absence of the apocrine component may have been additional contributing factors to the success of treatment in our patient.

In conclusion, ultrapulse CO<sub>2</sub> laser presents a promising modality for the ablation of NSJ in selected cases in cosmetically important areas. The lesions with the deep apocrine component, as well as involvement of the lower half of the dermis would qualify better for surgical excision than laser ablation because the potentially higher risk of recurrence with the later. Nevertheless, even if the cosmetic result is acceptable with the laser, and the lesion superficial, surgical excision may be a better choice if a thin-line scar can be achieved.

Although the treatment with the CO<sub>2</sub> laser was claimed to obscure early epidermal changes, with careful and prolonged follow-ups any malignant change could be recognized and treated quickly before the tumor achieves the significant size. Also the risk of malignant transformation of NSJ on the face is considered to be low<sup>3,4</sup>.

## REFERENCES

1. Mehregan AH, Pinkus H. Life history of organoid nevi. *Arch Dermatol* 1965; 91: 574-588.
2. Weng CJ, Tsai JC, Chen TJ. Jadassohn's nevus sebaceus of the head and neck. *Ann Plast Surg* 1990; 25: 100-102.
3. Wilson-Jones E, Heyl T. Nevus sebaceus. A report of 140 cases with special regard to the development of secondary malignant tumors. *Br J Dermatol* 1970; 82: 99-117.
4. Domingo J, Helwig EB. Malignant neoplasms associated with nevus sebaceous of Jadassohn. *J Am Acad Dermatol* 1979; 1:545-556.
5. Kauver ANB, Geronemus RG, Waldorf HA. Char-free tissue ablation: a comparative histopathological analysis of new carbon dioxide (CO<sub>2</sub>) laser systems (abstract). *Laser Surg Med* 1995; 16(Suppl 7): 50.
6. Alessi E, Sala F. Nevus sebaceus - a clinicopathologic study of its evolution. *Am J Dermatopathol* 1986; 8: 27-31.
7. Tarkhan II, Domingo J. Metastasizing eccrine porocarcinoma developing in sebaceous nevus of Jadassohn. *Arch Dermatol* 1985; 121: 413-415.
8. Baker BB, Imber RJ, Templer JW. Nevus sebaceus of Jadassohn. *Arch Otolaryngol* 1975; 101: 515-516.

9. Ashinoff R. Linear nevus sebaceus of Jadassohn treated with the carbon dioxide laser. *Pediatr Dermatol* 1993; 10: 189-191.

10. Ratz JL, Bailin PL, Wheeland RG. Carbon dioxide laser treatment of epidermal nevi. *J Dermatol Surg Oncol* 1986; 12: 567-570.

#### AUTHORS' ADDRESSES

Aleksandar L. Krunic MD, PhD, Director of Dermatologic Surgery, Dermatologic Surgery Unit, Department of Dermatology, University of Belgrade School of Medicine, Pasterova 2, 11000 Belgrade, Yugoslavia, and Dermatologic Surgery Unit, Division of Dermatology, Duke University Medical Center, Durham, North Carolina 27710, USA.

Robert E. Clark MD, PhD, Dermatologic Surgery Unit, Division of Dermatology, Duke University Medical Center, Durham, North Carolina 27710, USA

Mirjana M. Urošević MD, Department of Dermatology, University of Belgrade School of Medicine, Pasterova 2, 11000 Belgrade, Yugoslavia.