

# *Molecular aspects of skin ageing: recent data*

C. Rocquet and F. Bonté

---

## ABSTRACT

As our body's envelope, the skin acts as a biosensor with the environment and reflects our personality. Skin ageing is therefore an important and interesting topic of study. It results from the combination of intrinsic ageing and photoageing, which is due to the environmental influence, such as reactive oxygen species (ROS). The more recent data are gathered here to remind current knowledge about skin ageing, from a molecular level to the clinical signs, wrinkles and spots mainly. Because knowledge of the preferential biological targets of ageing has recently been making progress, it is possible to delay the manifestation of ageing, by acting on key biological processes.

---

## 1. Introduction

Humanity is ageing. The average life expectancy of people living in industrialized nations has doubled since 1900. This will result in social, economic, and health-care changes that will, in turn, drive public policy worldwide. The cosmetic industry is moving to cater for the ageing population by developing more innovative products. The skin provides a large interface with the environment, and is thus of prime importance. The major changes in the skin that occur with age are a loss of elasticity and a reduction in its protective function. These changes do not just affect the elderly, as they begin when people are younger than thirty. The extent of change depends to a large degree on how much the skin is exposed to sunlight and how stressful is a person's lifestyle.

For years, people have attempted to hide, reverse or control changes in their skin, such as wrinkling, roughness, mottling, blotching and dryness. Skin of elderly people is thin and fragile, due to complex changes very often summarized to reduced dermal collagen and decreased cell proliferation. Skin ageing is thought to result from two processes, intrinsic ageing and extrinsic, or photoageing (1, 2). Intrinsic ageing is believed to be genetically programmed and is thus presented as independent of all external and environmental influences. Extrinsic ageing is due to UV radiation and other environmental insults; that accelerate the skin changes (3). Photoageing leads to a rough/dry/leathery skin, a yellow/dull complexion, lentiginos or actinic spots and wrinkles. The study of the biology of ageing has made

## KEY WORDS

skin ageing,  
DNA, Asian  
skin, reac-  
tive oxygen  
species,  
wrinkle, cell  
senescence

rapid progress recently, after decades of stagnation. This paper reviews the most recent data on skin ageing, from the cell nucleus to tissue, and discusses their consequences for the skin.

## 2. Damage due to Reactive Oxygen Species (ROS)

Most people agree that free radicals are most important agents of ageing. Organic molecules absorb light or UV irradiation and become excited because an electron is transferred to a higher orbital. The excited molecule is a free radical which may cause secondary reactions and damage to various constitutive molecule (4). The energy storage molecule, adenosine triphosphate (ATP) is produced by oxidative phosphorylation in the mitochondria. The energy is produced by the oxidation of reducing equivalents of nutrients via the respiratory chain. However, mitochondria also regulate the intracellular calcium concentration and apoptosis. Abnormally increased membrane potential is linked to ROS (Reactive Oxygen Species) augmentation and mitochondrial DNA (mtDNA) mutations, constitutive to skin ageing.

ROS are normally produced by mitochondria, but they can also be produced following an external stress. Normal metabolism gives rise to most ROS, primarily via the mitochondrial respiratory chain, in which excess electrons are donated to molecular oxygen ( $O_2$ ) to generate a superoxide anion ( $O_2\bullet^-$ ). The superoxide anion is reduced by the enzyme superoxide dismutases (SOD) to hydrogen peroxide ( $H_2O_2$ ), which, in turn, is reduced to water by catalase, located in the primary peroxisomes, and by glutathione peroxidase (GPx), located in the mitochondria and cytosol. There are three isoforms of SOD (SOD 1 in the cytosol, SOD 2 in the mitochondria and SOD 3 in the extracellular space). Hydrogen peroxide can be converted to the highly toxic hydroxyl radical ( $OH\bullet$ ) in the presence of transition metals, and all three of the ROS ( $OH\bullet$ ,  $O_2\bullet^-$ ,  $H_2O_2$ ) can damage macromolecules directly or indirectly (5).

ROS are responsible for structural and functional alterations of cellular membranes, polyunsaturated fatty acids, proteins and DNA (6). For example, recent studies have shown that mitochondrial aconitase, an enzyme of the citric acid cycle that is critical for controlling the rate of ageing, is a target of oxidative damage (7). Cells also contain antioxidant enzymes such as superoxide dismutase, glutathione peroxidase, and catalase, and reducing agents like vitamin E and glutathione (8-10). Ageing is associated with a decrease in the plasma concentration of antioxidants, such as glutathione, and with increases in markers of oxidation damage, such as lipid peroxidation products (11,12). The total glutathione concentration in cultured skin fibroblasts decreases with

**Table 1. Reactive Oxygen species.**

FREE RADICALS	Half Life (at 37°C)
Hydroxyl	$OH\bullet$ 0.3 ns
Superoxide	$O_2\bullet^-$ 0.4 ns
Nitric oxide	$NO\bullet$ Seconds
Peroxynitrite	$ONOO(-)$ 50 ms – second
Singlet oxygen	$^1O_2$ $\mu$ s
Hydrogen peroxide	$H_2O_2$ Stable
Peroxide	$LOOH$ Stable

age, while glutathione reductase activity is unaffected. The antioxidant defenses are less well developed in the early stages of life than in postnatal life. (13).

The first changes that occurs in the skin after chronic or acute UV irradiation is the generation of reactive oxygen species, leading to the peroxidation of unsaturated lipids in the cell membrane. Low phototype individuals are more likely to produce large amounts of ROS after exposure to UVA, in particular singlet oxygen that can diffuse across the cell membrane. For example, the oxidation of catalase by singlet oxygen gives rise to more acidic conformers. The integrity of catalase may be a marker of the stress due to exposure to UVA (14). Singlet oxygen, due to its reactivity, appears more and more as a powerful free radical able to damage numerous skin components.

### *Mitochondria: an important target*

The mitochondrial theory of ageing states that mitochondria are the main site for generating free radicals and reactive oxygen in the cell. Thus, mitochondria are vulnerable to oxidative stress and damaged mitochondria can cause an energy crisis in the cell, leading to senescence and tissue ageing. An accumulation of damage decreases the cell's ability to generate ATP, so that cells, tissues, and individuals function less well. There is now considerable evidence that mitochondria are altered in the tissues of ageing individuals and that the damage to mtDNA increases 1000-fold with age. The phagocytic lysosome system for removing mitochondria is also considerably altered in the cells of ageing organisms. Thus, damaged mitochondria play an important role in apoptosis. The survival of the whole cell may depend on the release of caspase activators such as cytochrome C from the mitochondria (15). The extent of this alteration during senescence is still debated (16,17). Oxidative stress increases the production of free radicals and the inner membrane of the mitochondrion is chemically and physically altered. ROS are normally produced in mitochondria because of ubiquinones and the cytochrome b family (Complex III). Complexes I and IV are less vulnerable, while complexes II, III and V are affected by oxidative stress, via damage to lipids

and proteins (18). The electron transport chain is thus compromised, leading to energy supply failure and cell death (19). The mutation rate of mitochondrial DNA is ten-times higher than that of nuclear DNA. Indirect observations suggest there is transport of nucleic acids between mitochondria, which helps to repair the damaged mitochondrial genome (20). Damage to mtDNA will block mitochondrial turnover and replication, leading to decline in ATP production and protein synthesis. The accumulation of 8-hydroxydeoxy guanine (8-oxo-dG) in mitochondria also indicates great oxidative damage (21-23). Exposure of the human skin to solar radiation leads to an accumulation of mtDNA mutations, caused partly by oxidative damage, and these mutations play an important role in photoageing (24).

### *Genes and cellular pathways*

Alterations in oxidative metabolism and the cell redox state can affect many genes and cellular pathways. The influence of oxidation on mitogenic responses and signal transduction pathways, such as MAP kinase and NF- $\kappa$ B, are well documented (25). Human fibroblasts exposed to several oxidative stresses also develop markers of replicative senescence. Some genes, such as those encoding fibronectin, osteonectin,  $\alpha$ 1(I)-procollagen, apolipoprotein, SM22 (putative Ca binding protein overexpressed in senescent fibroblasts), and GTP- $\alpha$  binding protein are overexpressed. The mitogenic response to several growth stimuli (serum, PDGF, basic FGF and EGF) is lost (26). The reduction in connective tissue growth factor by UV radiation may contribute to the reduced procollagen synthesis observed in UV-irradiated normal human skin (27).

Peroxynitrite (ONOO $^-$ ) is generated from the transducer molecule nitric oxide (NO) and superoxide anions (O $_2^{\bullet-}$ ) under pathological conditions. MMPs and pro MMPs have been shown recently to be activated by peroxynitrite *in vitro* (28). The activation of poly(ADP-ribose) polymerase by peroxynitrite is also implicated in the pathogenesis of various inflammatory conditions and injuries. Tyrosine nitration is mostly mediated by peroxynitrite, a cytotoxic oxidant derived from nitric oxide that can cause DNA breaks. Peroxynitrite inhibits cell proliferation and high concentrations are also cytotoxic. Peroxynitrite and poly(ADP-ribose) polymerase also seem to be involved in the regulation of keratinocyte function and death in contact hypersensitivity (29). Nitrogen dioxide and carbonate radical anion must also be taken in consideration. Nitrogen dioxide can be produced from excessive nitric oxide autoxidation in hydrophobic environments such as cell membranes (nitrated lipids) and the interior of proteins (nitrated proteins); carbonate radical anions are produced in strong oxidative conditions and can oxidize nucleic acid guanine residues, GSH and proteins (30).

## **3. Proteins Oxidation and its consequences**

Thin, wrinkled skin is very often attributed to a lack of collagen. The dermis and overlying epidermis of ageing skin are profoundly altered (31). Slower protein synthesis is one of the most common events observed during ageing. The synthesis of both structural proteins, such as collagen, and enzymes that repair and maintain the normal metabolic functions of the cell, is slowed down. This leads to the inefficient removal of damaged molecules and decreased intra- and intercellular signaling pathways. The age-related increase in oxidized proteins may also be linked to modifications of proteins caused by lipid peroxidation products (32-35). Age-related changes in fibroblasts due to the metal-catalyzed oxidation of proteins lead to an exponential increase in the concentration of protein carbonyl groups in tissue samples taken from people aged 10 to 80 years.

Oxidative damage to proteins may be most important in ageing, because oxidized proteins become inactive and can accumulate in the cell, thereby triggering programmed cell death. ROS increase the carbonyl content of proteins by forming aldehydes and ketones from certain amino acid residues (36,37). The concentration of adenine nucleotide translocase, a protein in the inner mitochondrial membrane that is tightly bound to six molecules of cardiolipin, which contains highly unsaturated fatty acids, also decreases with age. This protein is the primary intracellular site for the generation of superoxide anion and exhibits adducts of the lipid peroxidation product, 4-hydroxynonenal, a powerful oxidative aldehyde. The concentration of cardiolipin may be a marker of the real age of the cell, based on its energetic capacity (38).

### *Proteasome involvement*

Protein oxidation *in vivo* is a natural consequence of aerobic life and the proteasome complex is responsible for the selective degradation of oxidized proteins. The age-related increase in the concentration of oxidized proteins is partly due to the cell's decreased capacity to degrade them. Lysosomal proteases and the proteasome complex normally degrade oxidized proteins. The 26S proteasome units selectively recognize and degrade oxidized proteins in the cytoplasm, the nucleus and the endoplasmic reticulum. The proteasome activity (multicatalytic proteinase MCP) involved in degrading oxidized proteins may be reduced (39). One of the lipid peroxidation products, 4-hydroxy-2-nonenal (HNE), can cross-link proteins via their lysine residues. The accumulation of oxidized protein, lipofuscin and/or ceroid pigments during ageing may be due to the changes produced in proteins by HNE and their subsequent inhibition of the proteasome unit

(proteasome 20S). This would lead to a vicious circle of cytotoxic protein oxidation products. (34). Oxygen stress (especially 4-hydroxy-2-nonenal) may attack proteins directly or through lipid peroxidation, to inhibit enzymatic activity. Oxidative damage to membrane transport proteins leads to alteration of the intercellular concentrations of calcium and potassium. The activity of the cytosolic proteasomal system also declines during the proliferative senescence of human fibroblasts (40). UVA and UVB irradiation both alter proteasome function in human keratinocytes (41, 42). While UV-induced skin damage is ameliorated by retinol and all-trans retinoic acid, UV irradiation blocks retinoid signaling in human skin through the ubiquitin/proteasome-mediated degradation of nuclear retinoid receptors (43, 44). In the future, it would be interesting to focus on the existence of preferential protein targets.

### Glycation

Cross-links can also form between proteins by coupling glucose carbonyl group to aminoacids such as lysine. These compounds called Advanced Glycation End products (AGEs) bind covalently to other proteins, and can cause extensive damage. The collagen lattice formed by cross-linked type I collagen is undeformable (unglycated collagen is fully compactible). Cross-linking collagen fibrils also alters the physical and mechanical properties of the extracellular matrix and changes the organization of the intracellular actin cytoskeleton (45). Glycated collagen may modify normal cell adhesion (46). As adhesion is a fundamental cell function, each alteration can damage cell behaviour (apoptosis, etc) and then change tissue homeostasis. The dermis and elastic fibre network become glycated in people over 35 years of age and solar irradiation appeared to enhance it (47). The fluorescence of epidermal tryptophan moieties and collagen cross-links in the dermal matrix can also be considered to be good *in vivo* markers of photoageing (48). While there are very high concentrations of antioxidant enzymes (catalase, SOD) in the epidermis, the concentrations are much lower in the dermis. Protein glycation and advanced glycation may be inhibited by antioxidant components (49). Photoaged skin has significantly reduced concentrations of antioxidant enzymes in the stratum corneum and the epidermis, while the concentration of oxidized proteins in the upper dermis is increased. Acute exposure to UV irradiation depletes the catalase activity in the skin and increases protein oxidation (50).

## 4. Dermal Matrix alteration

### Fibroblasts

The importance of cytokines, and immune cell ho-

meostasis for ageing and its clinical signs is still not clear. Normal concentrations of cytokines seem to be required for skin cell homeostasis. A disturbance leads to defects or wrinkle formation. Histological studies of chronically sun-exposed skin show that the dermis contains inflammatory infiltrates (51), mostly perivascular and perifollicular. Mast cells are more abundant in photodamaged skin than in normal skin. They synthesize and release mediators that modulate directly or indirectly extracellular matrix production and degradation (including TNF $\alpha$ , TGF $\beta$  and prostaglandin 2). They release proteases that can degrade the ECM or activate the proenzyme forms of metalloproteinases. Ultrastructural studies have also shown infiltration of the epidermis by macrophage/dendritic-like cells. These studies have recently been confirmed, showing more epidermal dendritic cells, but fewer Langerhans cells in sun-exposed skin (52).

The balance of cytokines in the skin is altered during ageing and fibroblasts become less responsive to growth factors or cytokines. Physiological ageing in human fibroblasts seems to be particularly associated with an altered response to interleukin-1- $\beta$ , a cytokine produced by monocytes, macrophages and other transitory cells involved in inflammation (53).

Transforming growth factor (TGF)- $\beta$ 1 is a cytokine involved in the differentiation of fibroblasts to myofibroblasts. These myofibroblasts are very important for dermal strength and may be responsible for the contraction of the dermis (54). It acts on fibroblast collagen, fibronectin, glycosaminoglycans, elastin production, all




Figure 1. Cellular consequences of stress on proteins.

of which are important for the mechanical properties of the skin. The expression of the TGF- $\beta$ 1 gene in epidermal keratinocytes does not decline with increasing cell age. Hence, TGF- $\beta$ 1 does not appear such as the message from epidermis to dermis affected by age (55).

Altered cytoskeleton function may play a key role in the age-related changes in several cell types, because it is involved in a variety of functions that are altered with ageing (immunological, endocrine and neurological changes) (56). The age-related changes in the cytoskeleton, due to its involvement in metabolic processes and cell surface receptors expression, can indicate defective signal transduction. Aged fibroblasts, which very often contract collagen gels poorly and do not migrate well, have a disordered actin microfilament cytoskeleton and a reduced  $\alpha$ -2- $\beta$ -integrin. (57).

Postmitotic and mitotic cells age differently. Postmitotic cells never divide, such as nerve, muscle and fat cells. Mitotic cells, such as keratinocytes and fibroblasts, divide. Replicative senescence is the process that limits the number of cell divisions. As senescent fibroblasts and keratinocytes accumulate with age in human skin, this could explain the deterioration of the appearance and properties of the skin. Senescent cells secrete degrading enzymes that modify the cytokine / interleukin balance, causing the loss of functional integrity (58).

Lipofuscin, or age pigment, accumulates in cells within the lysosomal vacuoles, especially in fibroblasts. Lipofuscin can also accelerate ageing and senescence under mild hyperoxia (59). The accumulation of lipofuscin may also be involved in spot formation. A lack of glucose-6-phosphate dehydrogenase (G6PD), an enzyme involved in the cell redox balance, may also accelerate fibroblast senescence (60). It has been proposed that phospholipid hydroperoxide glutathione peroxidase (PHGP) helps to protect fibroblasts against UVA-induced lipid peroxidation and the activation of metalloproteinase 1 (MMP1) by UVA.

### Receptors

The number, affinity and rate of internalization of epidermal growth factor (EGF) receptors are different in young and old fibroblasts, explaining the loss of responsiveness to EGF with age and the impaired wound healing in the elderly (61). Extracellular matrix is also diminished during ageing and the amount of collagenase in the skin increases with age. Collagenase production is controlled by protein kinase C via the members of the AP1 transcription factor family and can be inhibited by  $\alpha$ -tocopherol (62). Down-regulation of ligand-activated receptors is important for normal cell functioning. Receptors bearing their ligand move to specialized regions in the plasma membrane. The re-

sulting vesicles are transported through the cytoplasm by microtubules and fuse with endosomes and lysosomes, where they are degraded. ROS can also alter receptor function. For example, oxidase stress caused by hydrogen peroxide rapidly inhibits the internalization of receptor-bound EGF in human fibroblasts, so that the breakdown of the EGF-receptor complex is inhibited. Hydrogen peroxide also alters negative feed-back within the cell, and attenuates growth factor-induced signal transduction, leading to altered cell metabolism (63). Photo-damage increases the tenascin in cells, which might cause competition for the  $\alpha$ 2 $\beta$ 1-integrin receptor, reducing cell-collagen binding.  $\alpha$ 2 $\beta$ 1-integrin is the major collagen receptor, but is also a receptor for tenascin (64). Tenascin C is a large extracellular matrix glycoprotein whose production by keratinocytes is increased in wound repair; it is also found in normal adult skin. It is often distributed discontinuously in the upper papillary dermis adjacent to the EDJ, close to capillary basement membranes. The concentration of tenascin is increased in photodamaged skin and its distribution is different from that of skin protected from the sun. Tenascin is found along the dermo-epidermal junction in a continuous pattern and extends further into the papillary dermis. The pattern of gene expression in senescent fibroblasts is different from that of their still-dividing counterparts. Presenescent fibroblasts have low concentrations of metalloproteinases (MMPs) and high concentrations of MMP inhibitors (TIMP-1 and TIMP-3). The concentration of MMPs increases as cells become senescent, while that of TIMP decreases. The MMP production is stimulated by activation of the redox-regulated transcription factor NF $\kappa$ B and protein kinase C via activator protein 1 transcription factor (AP1) (65).




Figure 2. Cellular consequences of stress on receptors.

## MMP

Some of the MMP, a family of at least 16 enzymes that digest matrix macromolecules, are activated by UV irradiation (66). Thus, metalloproteinases 1, 3 and 9 (MMP-1, 3 and 9) in the epidermis are activated by UVB, while UVA stimulates MMP-1 *in vivo* and MMP-2 and 3 *in vitro*. The MMPs in the skin are responsible for breaking down macromolecules of the skin ECM, which ensures the skin's three-dimensional integrity. The balance between MMPs and MMP inhibitors is perturbed by environmental factors, such as light. This leads to collapse of the ECM and the visible effects of UV damage: wrinkling, loss of elasticity. Besides chronological ageing, actinic ageing, also called photodamage, causes premature skin ageing: thinning of the dermis, a loss of collagen content and protein organization and a breakdown of the ECM (67, 68).

Type 1 MMPs (interstitial collagenase) and type 9 MMP (gelatinase) break down skin collagen fibers, particularly during photodamage (69, 70). MMP-2 (gelatinase) acts on collagen types I, IV, and VII. Gelatin, elastin and fibronectin are all substrates for MMP-2, whose activity increases with age (71, 72). MMP-1 degrades collagen, which accounts for at least 70% of the dry weight of the dermis. Smoking increases the activity of MMP-1 in the skin *in vivo*. It leads to an imbalance between MMP-1 and the tissue inhibitor of metalloproteinase 1 (TIMP-1), which could be important for ageing (73). The MMP-1 produced by epidermal keratinocytes and dermal fibroblasts in response to various stimuli (74-78) appears to play a key role in dermal remodeling (79-81). Skin fibroblasts produce MMP-1 in response to UVB irradiation and keratinocytes play a major role through an indirect paracrine mechanism involving the release of epidermal cytokine after UVB-irradiation (82). MMP are produced in response to UVB irradiation *in vivo*, and are considered to be involved in the changes in connective tissue that occur in photoageing (83). They are associated with a variety of normal and pathological conditions that involve degradation and remodeling of the matrix (84-87). Several MMPs are produced during wound healing, such as MMP-3 in epidermis repair (88, 89).

UV rays and ageing lead to excess proteolytic activity that disturbs the skin's three-dimensional integrity. These proteinases are important for breaking down the extracellular matrix during chronic wound repair, in which there is re-epithelialization by keratinocyte migration (90). Thus, MMPs are continuously involved in the remodeling of the skin after chronic aggression.

Thrombospondin 2 (TSP2), a secreted extracellular matrix glycoprotein, is an adaptor and modulator of cell matrix interactions (91). It binds to heparan sulfate proteoglycan, low-density lipoprotein receptor-related protein (LRP), and the integrin  $\alpha\text{v}\beta\text{3}$  (92, 93). Increased MMP-2 activity (gelatinase A) leads to reduced fibro-

blast adhesion which could contribute to abnormal collagen fibril structure in the skin and the release of angiogenic factors. New data show that TSP2 binds to and inhibits MMP-2 indirectly and therefore plays a role in cell-matrix interactions (94).

The age modulated hypoxia response causes an imbalance between MMP-1 and MMP-9 and TIMP. Hence, there may well be altered MMP and TIMP gene expression at wrinkle sites (95, 96). Photodamage also results in the accumulation of abnormal elastin in the superficial dermis, and several MMPs have been implicated in this process. The quantities of matrilysin (MMP-7) and human macrophage metalloelastase (MMP-12), which have broad substrate specificities, are two key parameters that can be used to evaluate long term antiageing treatments (97). They are increased in the abnormal elastic fibers of chronically photoaged skin and contribute to the remodeling of elastic areas in sun-damaged skin. Human metalloelastase also aids macrophage migration, in addition to degrading elastic tissue, so amplifying the disturbance of the inflammatory homeostasis of the tissue. Ultrastructural and histopathologically studies have demonstrated that sun-exposed skin contains accumulated insoluble material and the normal elastic fiber architecture is lost, resulting in a loss of skin resilience and elasticity and probably wrinkle formation.

## Fibrillin and Elastin

Elastic fibers in the dermis form an amorphous matrix of elastin and intertwining bundles of microfibrils, which measure 10-14 nm in diameter. The oxytalan fibers are rich in microfibrils and are orientated perpendicularly to the basal lamina of the epidermis. A study on photoaged skin has shown that UV irradiation increases the tropoelastin mRNA in keratinocytes and fibroblasts (98). Selective inhibition of skin fibroblast elastase could be one way to fight wrinkle formation following cumulative ultraviolet B irradiation (99). Lysozyme may alter the elastic fibers in the surface, preventing further degradation and the accumulation of altered elastic fibers.

Photoaged skin contains elastic material in the reticular dermis, and the fibrillin deposits in the reticular dermis are enlarged. Elastic fibers have a central core of hydrophobic cross-linked elastin surrounded by fibrillin-rich microfibrils. The papillary dermal microfibrillin-rich microfibril network is truncated and depleted in photoaged skin. There are fewer fibrillin-rich microfibrils in wrinkled photoaged skin, probably due to inflammatory cell proteinases (neutrophil elastase), or activation of matrix metalloproteinase (100). Cross-linking causing decreased elasticity could also be involved in wrinkle formation (101).

The fluorescence of tryptophan and collagen cross-links in the dermal matrix may serve as *in vivo* markers

Table 2. Main Matrix Metalloproteinases and their substrates.

Name	Enzyme	Selected substrates
<b>MMP-1</b>	Matrix collagenase (fibroblast collagenase)	Collagens I, II, III, VII and X
<b>MMP-8</b>	Neutrophil collagenase	Collagens I, II, III, Link protein, Aggrecan
<b>MMP-13</b>	Collagenase 3	Collagens I, II, III
<b>MMP-18</b>	Collagenase 4	Collagens I
<b>MMP-2</b>	Gelatinase A	Gelatins, Collagens I, IV, VII and XI, Fibronectin, Laminin, Elastin
<b>MMP-9</b>	Gelatinase B	Gelatins, Collagens IV, V and XIV Aggrecan, Elastin
<b>MMP-3</b>	Stromelysin 1	Aggrecan, Gelatin, Fibronectin, Laminin, Collagens III, IV, IX and X
<b>MMP-10</b>	Stromelysin 2	Aggrecan
<b>MMP-14</b>	(membrane type)	Collagens I, II and III, Laminin
<b>MMP-7</b>	Matrilysin	Aggrecan, Fibronectin
<b>MMP-11</b>	Stromelysin 3	Fibronectin
<b>MMP-12</b>	Metalloelastase (Macrophage)	Elastin

of skin aging, photoaging, and as a way of assessing exposure to UVA radiation (48). The morphology of elastic fibers changes significantly during life. The number of elastin microfibrils (mainly composed of fibrillin) gradually decreases during ageing, and the degenerative process is accelerated by exposure to sunlight. Amyloid P and lysozyme are deposited in thickened fibers, while amyloid P alone is deposited in oxytalan vertically oriented fibers in the papillary dermis. Deep wrinkles are linked to the degeneration of collagen and the deposition of abnormal elastic material (102). Wrinkles are formed by major changes in the dermis matrix and at the dermo-epidermal junction. The content of fibrillin, a component of the elastic fiber network, is increased by prolonged clinical doses of topical retinoic acid. This is why fibrillin-1 has been proposed as a "reporter" molecule for the efficacy of photoaging.

#### *Water and GAG*

The altered skin texture and structure of elderly people is caused by changes in proteins, lipids and water, leading to altered mechanical properties, such as wrinkling, sagging, loss of elasticity and apparent dryness. Water structure is important because water can bind to various proteins and is important for maintaining the structural and mechanical properties of proteins. Their natural interaction is diminished in photoaged skin, leading to decreased collagen stability and the fragmentation of collagen fibrils (103). The distribution of glycosaminoglycans (GAG) in the dermis seems to be modified in sun-damaged skin and could be linked to alterations of deep protein. Studies using immunoperoxidase staining of hyaluronic acid and chondroitin sulfate and confocal laser scanning microscopy have

shown increased dermal GAGs in sun-damaged skin. The GAGs are deposited on the elastic material of the superficial dermis and not between collagen and elastic fibers, as in normal skin (104). Hyaluronan is a major constituent of the skin extracellular matrix. Hyaluronan polymers become more tissue-associated with advancing age (105). Together with changes in proteins, this contributes to the pronounced alteration of the skin mechanical properties in the elderly.

#### *Lipids and cell membrane*

The skin barrier is linked to the lipids of the intercorneocyte space. Intercellular lipids consist of an organized mixture of ceramides, sterols and fatty acids (106, 107). The lipids in intercellular membranes form short- and long-periodicity lamellar phases (108). A recent X-ray diffraction and electron microscopy study showed no correlation between differences in the organization of stratum corneum lipids and ageing, despite the changes in skin properties often observed in the elderly (109).

Vitamin A (retinol and retinyl esters) is present in the epidermis as free and esterified retinol. Acute exposure to UVA completely depletes the epidermis of vitamin A and causes lipid peroxidation. In contrast, exposure to UVB results only in the loss of vitamin A (110).

The human sebaceous gland undergoes both extrinsic and intrinsic ageing (morphological changes in the sebaceous gland activity). The highly androgen-dependent sebum secretion in neonates reaches its maximum in young adults. The number of sebaceous glands remains unchanged throughout life, but sebum production tends to decrease after menopause in women and after the eighth decade in men. The age-dependent

decrease in androgen leads to a slower cell turnover in the sebaceous glands, resulting in hyperplasia of the facial sebaceous glands. UV may contribute to this process. Molecular studies have shown that overexpression of the ageing-associated gene *Smad7* and parathormone-related protein are linked to hyperplasia of the sebaceous gland, but overexpression of the *c-myc* gene is associated with enhanced sebum production (111). Decreased sebum production is also responsible for skin dryness in the elderly.

## 5. DNA Damages and consequences

The DNA of the skin is constantly submitted to environmental damage and has developed mechanisms to repair this damage. The balance between damage and repair has a major impact on ageing. The most common cause of premature skin ageing is UV irradiation, which damages DNA through photoproducts. DNA repair is essential for maintaining the functional integrity of DNA. Selective repair, such as the removal of pyrimidine dimers, occurs in the transcribed strand. DNA is repaired by a variety of mechanisms, such as direct reversal DNA damage for thymine dimers, base excision and mismatch repair. Nucleotide excision repair is very important for damage caused by UV. A repair complex binds the damaged DNA, then an endonuclease cuts it on either side of the damaged nucleotide. The original DNA sequence is resynthesized by a DNA polymerase and a ligase.

The most reactive oxygen free radical,  $\text{OH}\cdot$ , reacts with DNA bases to give altered bases, such as 8-hydroxydeoxy guanine. These are eliminated by the DNA repair enzyme complex, but some accumulate with ageing (112). Ageing may also result from the injury to mitochondrial DNA and peroxidation of the inner mitochondrial membrane lipid. Many studies have suggested that mtDNA suffers more from oxidative DNA damage than does nuclear DNA.

Genomic and mitochondrial DNA are both intimately involved in the process of ageing. There is some decline in DNA repair capacity with age. Increased DNA fragility or DNA strand breaks, chromosomal aberrations based on cytogenetic examination, decreased DNA methylation and changes in ploidy all increase with age. Rearrangements, translocations, and sequence alterations also increase with age (113). The main type of damage generated by apparently all types of reactive oxygen species (ROS) is oxidative changes to guanine (114).

Free radicals produce a number of lesions in DNA, damaging bases, sugar lesions, DNA-protein cross-links, causing single-strand breaks, double-strand breaks, and abasic sites by different mechanisms (115). A recent

review relates the contributions of stress-induced damage to cellular DNA:

- by damage to nuclear DNA and its repair mediated by poly(ADP-ribose) polymerase1 (PARP 1)
- damage to telomeric DNA and its contribution to telomere-driven cell senescence
- the accumulation of mutations in mitochondrial DNA (116).

The effects of oxidative stress can be direct or indirect. For example, some cell constituents (flavins, porphyrin), many dyes (acridines, methylene blue, neutral red) and drugs can act as photosensitizers inside cells. The excited state of a photosensitizer can be thought of as genotoxic species similar to other free radicals, because they can directly or indirectly cause DNA modifications (4). Highly selective changes to guanine are also caused by photosensitizers that modify DNA via singlet oxygen (type II photoreaction), or one-electron oxidation (type I photoreaction) (117-120). Endogenous and exogenous oxidative stress can cause serious damages to mitochondrial DNA. Deletions of mitochondrial DNA may be used as markers of skin ageing and exposure to UV irradiation (121-124). Direct evidence for the increased presence of UV-induced damage in mtDNA was obtained recently. Ray *et al* (2000) used a PCR method to show that the number of mtDNA deletions in the epidermis is significantly associated with increased exposure to UV radiation. UV radiation may directly or indirectly act via free radicals to cause mutations at labile sites in mtDNA, enhancing intra genome recombination, and increasing deletions. Mutations of mtDNA accumulate during ageing and in photoaged skin; the most common mutation is a 4977 base pair deletion (called common deletion). Chronic exposure of human skin to sunlight results in more mtDNA mutations than in un-exposed skin. UVA-irradiation produces singlet oxygen that generates the common mutations of mitochondrial DNA that occur in photoageing (125).

It is difficult to precisely measure oxidative DNA damage, because extraction and sample treatment may cause oxidation. Various analytical techniques can be used to measure oxidative damage to DNA: gas chromatography (GC) and liquid chromatography (LC) with mass spectrometry (MS) provide positive identification and accurate quantification. Modified nucleosides have been measured recently by methods using LC/tandem MS (LC/MS/MS) and LC/MS (112).

## 6. Telomerase involvement

The closest thing to a cellular clock resides at the tips of chromosomes. The chromosome ends, the telomeres, do not contain genes that program hereditary traits, but are functional complexes. The telomeres at the ends of eukaryote chromosomes protect them from



degradation or fusion. The telomeres shorten each time human cells divide. The cell may finally fall into a senescent state. Telomerase is a ribonucleoprotein that synthesizes the repeated sequences at chromosome ends and helps DNA polymerases to complete the replication. New data indicate a new way to link telomeres to senescence. The telomere is a dynamic nucleoprotein complex that can switch stochastically between an uncapped state and a capped state, which preserves the physical integrity of the telomere and allows cell division to proceed (126-130).

It has been suggested that cell senescence may be good because it is a defense against cancer, which is marked by uncontrolled cell division. Cells unable to regrow their telomeres stop dividing before they can cause too many mistakes. As there is no telomerase in many somatic tissues, telomere erosion may well be a major factor in cell ageing (131). Telomerase therapy might one day help generate a new supply of cells to treat age-related diseases. It has been showed that there is a telomerase activity in the skin.

It is possible to distinguish between ageing and longevity. Telomere shortening is probably more involved in regulating cell longevity than in ageing *stricto sensu*. Ageing is due to accumulation of molecular disorders and a lack of cell energy. This loss of energy is involved in the deterioration and a gradual loss of the functional integrity of the tissues. Nevertheless, telomere shortening (or more precisely telomerase dysfunction), oxidative damage and hormones could all be signals involved in ageing.

The regulation of telomerase in mammalian cells is multifactorial, involving telomerase gene expression, post-translational protein-protein interactions, and protein phosphorylation. Several proto-oncogenes and tumor suppressor genes are involved in the regulation of telomerase activity (132). Several physiological factors, like EGF and/or amphiregulin, and growth factors, can also influence telomerase (133). Telomere length and telomerase activity may determine cell senescence. Hyperoxia accelerates telomere shortening by causing oxidative stress. A reduction of stress, for example by the action of free radical scavengers, delays replicative senescence. Telomere act as a "sentinel" for oxidative damage to the genome and replicative senescence may be triggered by telomeres as a consequence of DNA damage. It is thus very important to ensure that cells have sufficient telomerase (134). Guanine of the telomerase 3' overhang (TTAGGG) can be considered as a target for reactive oxygen species or UV irradiation (135). Estrogen activates telomerase via direct and indirect effects. There may be hormonal control of telomerase activity. Sex steroids may thus influence cell senescence and ageing (136). The activity of telomerase may be also regulated by the tumor suppressor protein p53; a lack of this protein may lead to increased telomerase activity in cancer cell development (137). Mutations of the p53 gene and telomerase activity are linked, and these mutations are considered to be UV specific (138). The transcription factor NF- $\kappa$ B may act at a specific site to influence the activity of the telomerase catalytic subunit (hTERT) T (139). More recently,




Figure 3. Consequences of external stress on the DNA.

telomerase and its catalytic subunit hTERT have been shown to be involved in oxidative stress, and UV irradiation disturbs the telomerase activity in human keratinocytes (140).

The nuclear enzyme poly(ADP-ribose) polymerase 1 (PARP 1) participates in the regulation of both DNA repair and transcription. Moderation of PARP following DNA damage has also been proposed to protect skin cells from UV induced acute and chronic photodamage (141). The ageing and survival of endothelial cells are linked to molecular mechanisms that control cell proliferation, quiescence, apoptosis and senescence. The activation of telomerase in human dermal microvascular endothelial cells also seems to affect their durability both *in vitro* and *in vivo* (142).

### Apoptosis

Cell senescence could be, like apoptosis, a part of the body's defenses, a natural mechanism to prevent cells from accumulating mutations with their physiological consequence, malignancy (143). The differentiation, apoptosis and senescence of keratinocytes share some molecular pathways. Epidermal differentiation and apoptosis lead to cell death and the removal of cells by transglutaminase activation and proteolysis. Regulators of apoptosis, such as Bcl-2 (suppressor) or Bax (promotor), are also produced during differentiation. Senescent cells are cells which can no longer replicate, but can still respond to growth factors. They have lost their ability to perform correctly tissue homeostasis. Cells lose their ability to divide early in differentiation.

Some authors think that senescence is not directly linked to epidermal differentiation, due to differences in the responses to cell cycle inhibitors (144).

Apoptosis is a cellular end point of the stress response. Apoptosis removes damaged cells from UV-irradiated tissues. The genes that control apoptosis in the epidermis, such as the bcl-2 gene, are disregulated during ageing. The decreased efficiency of apoptosis may contribute to chronological ageing and extrinsic skin ageing. Only epidermal stem cells escape cellular senescence. It appears that epidermal terminal differentiation, apoptosis and cell senescence are all triggered by stimuli. Nevertheless, keratinocytes will undergo classical epidermal differentiation or will irreversibly enter into senescence or apoptosis, depending on their state and on the nature and strength of the stimuli. Oxidative damage is a cellular stress that can cause senescence like growth arrest, or even apoptosis. Ras-induced senescence is also mediated by ROS, but is not clearly associated with telomere shortening. The tumor suppressor P16 accumulates as fibroblasts approach senescence, and by inhibiting its degradation, P19 indirectly mediates the growth arrest or apoptosis. P (145).

## 7. Morphological Variations

The skin becomes thicker until maturity and then becomes thinner in women over 50-60 years old. Measurements of skin physical properties show that it becomes thinner, stiffer, less tense and elastic with ageing (146).

### Mechanical Properties

Young's modules (elasticity modules) of the skin, a ratio between stress and deformation, increases linearly with age. This is in agreement with data indicating that skin becomes more rigid and less able to stretch in response to stress with age. This has to be correlated with the increased crosslinking of collagen, the disorganization of the fibril network and the large amount of free water in the dermis. Ageing decreases skin function and causes clinical changes such as wrinkling, color changes (yellowish, patches, pigmentation), and a loss of elasticity (146). Sagging is one of the major age-related morphological changes in the face. While wrinkles and general changes in the face have been well studied, a recent method using photostandards and 3 D analysis of replicas shows that women's cheeks begin to sagging when they reach 40 (147). Recently measurements of site-related and age-dependent variations in facial skin show that there is an overall increase of skin echogenicity and thickness with age. The skin on the upper and lower lips, on the chin and infraorbital regions is thicker than that on the central forehead, lateral forehead and cheeks. The facial skin thickness be-




Figure 4. Consequences of ageing on main biological targets.



**Figure 5. Clinical signs of ageing on a Caucasian woman of 75 years old: presence of wrinkles, spots and sagging.**


comes greater over the lateral regions of the forehead, lips and nose in elderly subjects, and becomes thinner over the infraorbital regions (148). Fine lines are due to the gradual breakdown of collagen and elastin fibers, and they are exacerbated by sun damage. Very deep wrinkles are associated with the muscle below the skin surface. Muscles contract more with age to compensate for the loss of volume. Excessive exposure to sunlight and smoking can cause major changes in the skin (73). The skin may darken; develop very fine wrinkles, spots, and sag, all of which are symptoms of photoageing. This is a very serious concern for middle-aged women, especially women in Asia. Studies using the two point gap discrimination method plus microneurographic recording in response to mechanical stimuli have also revealed changes in tactile spatial discrimination in the elderly (149).

The subcutis is also concerned in ageing. Ageing results in larger fat cells in the subcutaneous tissue. Hormone changes linked to ageing may also cause difference in body fat distribution (150). Energy metabolism is also regulated via leptin, a fat cells-derived hormone, in adults. The concentration of leptin in the blood varies during the menstrual cycle. Leptin binding activity

is low at birth and high in the pre-pubertal years, but it is stable during adult life and does not vary with ageing (151-153). The adipocytes also act as estradiol stores. The circulating concentration of this hormone varies with age, and is most important in mature skin. Menopause, the physiological cessation of menstruation caused by decreased function of the ovaries, leads to thinning of the dermis, mainly due to a decrease in the collagen content, atrophy of subcutaneous tissues and increased skin dryness (154, 155).

### Wrinkles

Wrinkles are modifications of the skin associated with cutaneous ageing and develop preferentially on sun-exposed skin. Clinicopathological features of wrinkles were studied among the different types of skin relief modifications. Four types of skin depressions can be defined according to their depth: folds, permanent wrinkles, reducible wrinkles and skin micro-relief. Development of wrinkles may be secondary to actinic elastosis and to the disappearance of microfibrils and collagen fibers at the dermoepidermal junction. Epidermis is involved with a decrease of the cell renewal, an increase of involucrin, a decrease of integrin  $\beta$  1, type VII collagen and fibrillin 1 (156). Wrinkles are one of the major concerns for women along with spots and freckles. Principal causes of wrinkling are ageing and excessive exposure to UV rays. Wrinkles are the expression of the accumulation of modifications at different levels of the skin. Development of so-called fine wrinkles begins to take place in the thirties, reaching a peak in the fifties, while deep wrinkles increase in the fifties. Little is known about the exact histological changes underlying wrinkle formation. Changes in collagen type I, III, type IV and VII at the DEJ have been recorded (157-160). Collagen VI, concentrated in the



**Figure 6.  $^{14}\text{C}$  Ascorbic acid intracellular transport in Human Normal Fibroblasts (lifting),  $n=12$  (M. Dumas courtesy).**

papillary dermis immediately below the dermal-epidermal junction is similar in photoprotected and photoaged skin (159). Some fibroblasts which are accumulating damages are less stimulated by surrounding ascorbic acid, resulting in a decrease of collagen and a loss of dermis density. Very recently, in aged fibroblasts from photoexposed zones, a decrease of the intracellular transport of pericellular ascorbic acid has been proposed in wrinkle formation and ellagic acid derivatives have been proposed to overturn the phenomenon (161; Marc Dumas, personal communication).

The balance between the stress activated (SAPK) and mitogen activated (ERK) MAP kinase signaling pathways regulates cell growth and extracellular matrix production (ECM). SAPK activity, measured by c-jun phosphorylation, is increased in the elderly and inhibits ECM production by activating collagenase and inhibiting collagen synthesis (162). Oxidative damage is central to skin ageing, and is particularly involved in wrinkle formation. The extracellular signal-regulated MAP Kinase pathway (ERK), which mediates cell responses to growth factors, is less active in old human skin *in vivo*. At the same time, the activity of the stress-activated MAP Kinase pathway (c-jun, p38 MAP Kinase) increases in old human skin *in vivo*. The amounts of c-Jun mRNA and protein are increased in old skin, but the amounts of c-Fos mRNA and protein are not. It has also been demonstrated that retinol activates the ERK pathway in old skin but does not alter the stress-activated c-jun kinase. (163). TGF  $\beta$ -1 in the extracellular matrix and in keratinocyte may also be an important marker for measuring the efficiency of an "anti-wrinkle" treatment (164).

### Spots / Freckles

Melanocytes are specialized cells that are located in the basal layer of the epidermis. They synthesize and transfer melanin pigments to surrounding keratinocytes, thus protecting from UV carcinogenic effects (160, 165, 166). Molecular mechanisms and cell cycle regulatory gene expression leading to melanocyte senescence and transformation differ significantly from fibroblasts. As found in other cell types, progressive telomere shortening appears to trigger replicative senescence in normal melanocytes. In melanocytes and not fibroblasts, there is a loss of p21<sup>Waf-1</sup> and cyclin E expression. Melanocytes and fibroblasts present common events as an increase in p16<sup>INK4a</sup> levels and down-regulation of E2F1, also shared by senescent keratinocytes. In fibroblasts, the senescent phenotype is linked to the repression of the gene c-fos, upregulation of p21<sup>Waf-1</sup>, and down-regulation of E2F transcription factors. p21<sup>Waf-1</sup> may be a major regulator of fibroblast senescence. (167).

Adult melanocytes are able to stop to proliferate and terminally differentiated melanocytes are still metabolically active but postmitotic. The altered differentiated functions of senescent melanocyte are not well known.

Alterations of melanosomes, melanin synthesizing enzyme in mitogen activated protein kinase (MAPK), and in cell cycle progression have been reported. All these altered functions may have a real impact in tissue (168). The knowledge of the mechanisms of human skin color is of prime importance to develop skin care increasing the skin radiance and fighting spot formation. The change of the absorbance spectrum from reflectance including the scattering effect has been found not to correspond to the molar absorption spectrum of melanin and blood (169, 170).

Disturbed keratinocytes-melanocytes interactions during melanosome transfer and skin melanosome distribution patterns could be related to spot formation. Melanocytes located in the basal layer of epidermis produce melanin-loaded melanosomes, which are distributed to neighboring keratinocytes. UV radiation lead to an accumulation of melanosomes in melanocytes and treatment with MSH induces exocytosis of melanosomes. UV and MSH increase the phagocytosis of melanosomes by keratinocytes (171).

Kinesin and kinectin are motor proteins that are involved in some stage of melanosome transport (166; 172-174). Melanosome transport along the dendrite is mediated in part by microtubule proteins and myosin V, which traps melanosomes at the actin-rich periphery of the dendrite SNARE protein (soluble N-ethylmaleimide-sensitive factor attachment protein receptors) and Rab-3a. Proteins involved in vesicle transport, docking (rab 3) and membrane fusion (SNARE) seem to be involved in the changed melanosome dynamics caused by UV irradiation. This can be thought of as an exocytosis towards the keratinocytes (175). Serine protease inhibitors that interfere with PAR 2 (protease-activated receptor 2) may cause depigmentation by affecting melanosome transfer and distribution (176). The formation of skin spots on photoexposed areas is a very complex problem, but may be viewed as a double problem. The direct and indirect processes stimulated by UV irradiation (endothelin-1, thymine dimers, NO) increase the melanin production (177-179). Their local aggregation can also generate new reactive oxygen species (180). A disturbance of cell-cell signaling (cadherin E, P, etc) can cause melanin to be transferred to basal keratinocytes instead to suprabasal cells, resulting in pigment accumulation (181). Accumulation of pigment in such cells could also be due to local defect of keratinization and to oxidation products, like lipofuscin. Changes due to UV irradiation and ageing could make these cells, pigment-loaded by error, less easy to eliminate. Furthermore, melanocytes getting the information that their protective pigment distribution pattern is abnormal will continue to oversynthesize pigment, leading to more, uncontrolled accumulation.

Other biochemical factors may be responsible for spot formation. Melanogenic stimulatory factors derived from epidermal cells in senile freckles include endo-

thelin and stem cell factor (182). Abnormal sphingomyelinase production leads to high concentrations of sphingosylphosphorylcholine instead of ceramide in the epidermis of patients with atopic dermatitis. The sphingosylphosphorylcholine is associated with the pigmentation defects frequently observed in atopic dermatitis (183). The question of a specific way of spot formation in mature skin linked to the hormonal balance modification is also still open. UV-induced melanogenesis is mediated by nitric oxide radicals. There is a decrease in tyrosinase activity stimulated prior to NO-stimulation (184). Human melanocytes contain the mammalian melanin concentrating hormone (MCH), but human keratinocytes and fibroblasts do not. The MCH is coupled to a G protein receptor, SLC1; the inactivated complex blocks the cyclic AMP second messenger pathway and increases intracellular calcium. Disturbance of this pathway by UV irradiation or other elements can also favor spot formation (185). The hormonal control of skin pigmentation via interaction of POMC peptides (eg  $\alpha$ -MSH, ACTH) with other local and circulating hormones may influence melanocyte function, and thus coordinate the pigmentary response of the skin, especially after major changes in hormone concentration (186). Metallothioneine, an intracellular free radical scavenger, could be induced in human melanocytes. Suppression of melanogenesis is partly due to the induction of metallothioneine.

### *Cutaneous microvasculature*

The ageing and survival of endothelial cells are linked to molecular mechanisms controlling cell proliferation, quiescence, apoptosis and cellular senescence. The activation of telomerase in human dermal microvascular endothelial cells is linked to their durability both *in vitro* and *in vivo*. Knowledge of telomerase activity and other markers of amplifying dermal perivascular cells may reveal more about the regenerative capacity of the skin microvasculature. Telomerase activity / length seems to be directly linked to the angiogenic potential (130; 142; 187-190).

## *8. Specificity of Asian Skin*

The formation of facial wrinkles is a sign of photoageing. A unique study of over 3000 people from five ethnic groups (African, American, East Asian, Caucasian, Indian Asian and Latino) of different ages has revealed age-dependent changes of the skin (wrinkles, hyperpigmentation and pores). The mean fraction of the face area covered with wrinkles is significantly smaller in African Americans than in Caucasians, but East Asians have the smallest wrinkled area at any given age. The authors suggest that racial differences in other genetic factors besides skin pigmentation, such as DNA

repair, are important in determining the development of skin wrinkles. African Americans have more hyperpigmented spots and facial pores than other racial groups. Caucasians have significantly less well hydrated skin than African Americans, East Asians and Latinos (191). The relationship between skin phototype and deep and fine wrinkle scores on the faces of 230 Japanese subjects shows that sunlight-sensitive subjects have deeper wrinkles (192).

A recent study has compared the action of sun protection factor according to COLIPA recommendations on Asian and Caucasian volunteers. The observed difference in SPF is not only due to skin color but also to internal factors affecting response of the skin to UV (193). Asian skins are not all the same. Differences in the minimal dose causing erythema in Chinese and Korean subjects were also linked to differences in skin chromophores (194). A study of 230 Japanese individuals classified according to their skin phototype (type IV predominant) suggests that deep wrinkles are more severe in phototype I, and that there is no link between phototype and fine wrinkles. Skin phototype does not seem to be related to hair or eye color. Some Japanese with dark hair and black eyes have sun sensitive low phototype (192).

A study of the factors causing dark circles around the eyes in 60 healthy Japanese women indicates that the dark circles become darker as the blood mass increases, showing the importance of hemodynamics in the area (195). A comparison of the cheek skin color of Caucasian and Japanese women shows an increase in the yellow axis with age in Japanese women, whereas there was an increase in the red axis in Caucasians, followed by a decrease around 50 years (196). East Asians living in Los Angeles and Japanese living in Akita (Japan) had similar degrees of skin wrinkling. East Asians have less facial hyperpigmentation than Latinos, African Americans or Caucasians (191). The facial hyperpigmentation of 56 women (aged 22 to 67 years) and melanin spot distribution showed more melanin granules around the nuclei in chloasma and senile pigmentation (197). Age-related alterations in the echogenicity of the skin in Japanese women (130 women aged 18-83 years) were studied, with measurements on the forehead, eye corners, and cheeks. An increase in the lower layer of the dermis probably due to the accumulation of degraded or disarranged collagen and a decrease in echogenicity in the upper dermal layer indicate a tendency similar to that seen in Caucasians (198). Measurements of the sebum excretion rate and skin surface contours of 662 healthy Korean volunteers revealed age-related changes. The forehead/cheek sebum excretion rates increased with age while there were changes in the skin pores and skin surface texture. The disappearance of primary lines, secondary lines and increased pore size with age is probably linked more to exposure to sunlight (199). As demonstrated for the Japanese

volunteers, sunlight also modifies lipid peroxidation, and cholesterol 7-hydroperoxide is a good marker (200).

According to JH Chung, the patterns of wrinkles in Asian people differ from those of Caucasians. For example, Korean people have deeply wrinkled forehead and perioral areas. Wrinkles and pigmentary modifications are the two main characteristics of photoageing in Koreans (201). The skin of groups of 100 different Asian women was compared. In Japanese the skin was in a better condition compared to six other populations. Age-related increase of sallowness was more prominent in Chinese and Korean skin, while, in Philippines the highest tendency of combined skin lesions prevailed (202).

There are several independent risk factors of wrinkles, such as age, exposure to sunlight, menopause, skin color and smoking. Pregnancy is another factor of risk for facial wrinkling due to the high levels of sex hormone bound to globulin and low concentrations of free estradiol. Progesterone also acts as an antagonist to estrogen. Lactation delays the return of ovulation and decreases the estradiol concentration. Hormone Replacement Therapy (HRT) decreases the risks of wrinkling. The decrease in skin collagen due to estrogen deficiency in post-menopausal women may aggravate the severity of wrinkles. Men are also subject to pigmentation damage, leading to seborrheic keratosis and solar lentigo, and, as in women, it leads to depigmentation (lentigo, freckles, mottled pigmentation). In 189 Korean women, a significant increase in the risk of wrinkles was found associated with an increasing number of full-term pregnancies and menopausal age (203).

A multicentric study carried out on 3 000 Chinese women showed no crow's feet area prevalence in Northern cities and a higher peri-oral and glabella wrinkle score in Southern cities. Only 203 women were concerned at 21-25 years of wrinkles in crow's feet area, but at 36 years 75 % of women were concerned (204). Comparing 160 French and Chinese women (20-65 years) the onset of wrinkles delayed by around 10 years in Chinese women (205). From 26 to 60 years, 60% of the Chinese women exhibit pigmented spots on their face. Small facial pigmented spots characterize young population (18-40 years), spots of more than 6 mm diameter increase after 30. Whatever the age group, pigmented spots are always more pronounced on the face than on the hands. Climatic factors and chronic exposure of UV are suggested as a main cause of pigmented spots (206).

## Conclusion

The skin acts as a biosensor because it forms a large interface with our environment. It is a dynamic living barrier which is of prime importance in our social life. The recent studies described in this paper clearly show the complex interconnections at all levels of the tissue and provide a more precise picture of the importance of the events involved in ageing and photo-damage. It may become possible to delay the appearance of signs of skin ageing by acting on key mechanisms revealed by research into this fascinating topic

## Abbreviations

AGE	Advanced Glycation products
ADP	Adenosine di Phosphate
ATP	Adenosine tri Phosphate
DNA	Desoxyribonucleic acid
ECM	Extracellular Matrix
EDJ	Epidermal-Dermal Junction
EGF	Epidermal Growth Factor
FGF	Fibroblast Growth Factor
G6PD	Glucose-6-Phosphate Deshydrogenase
GAG	Glycosaminoglycans
GC	Gas Chromatography
GPx	Glutathion Peroxydase
GTP	Guanosine Triphosphate
HME	Human Metalloelastase
HNE	4-hydroxy-2-nonenal
HRT	Hormone Replacemnt Therapy
hTERT	telomerase catalytic subunit
LC	Liquid Chromatography
LRP	Lipoprotein receptor-related protein
MAPK	Mitogen Activated Protein Kinase
MCP	Multicatalytic Proteinase
MMP	Matrix Metalloproteinase
MSH	Melanocyte stimulating Hormon
mt DNA	Mitochondrial DNA
PARP	Poly(ADP-ribose) polymerase 1
PDGF	Platelet-derived growth factor
ROS	Reactive Oxygen Species
SAPK	Stress Activated MAP Kinase
SNARE	Soluble N- ethylmaleimide-sensitive factor attachment protein receptors
TSP2	Thrombospondin 2
UV	Ultraviolet

## REFERENCES

1. Trautinger F. Mechanisms of photodamage of the skin and its functional consequences for skin ageing. *Clin Exp Dermatol*, 2001; 26: 573-7.
2. Ma W, Wlaschek M, Tantcheva-Poor I, Schneider LA, Naderi L, Razi-Wolf Z, Schüller J, Scharffetter-Kochanek K. Chronological ageing and photoageing of the fibroblasts and the dermal connective tissue. *Clin Exp Dermatol*, 2001; 26: 592-9.

3. Podda M, Grundmann-Kollmann. Low molecular weight antioxidants and their role in skin ageing. *Clin Exp Dermatol*, 2001; 26: 578-82.
4. Epe B. DNA damage induced by photosensitizers and other photoreactive compounds. In: DNA and free radicals: Techniques, Mechanisms & Applications. OICA Internat. St Lucia, London, 1998; 1-20.
5. Wallace DC, Melov S. Radicals aging, *Nature Genetics*, 1998; 19: 105-10.
6. Benzi G, Pastoris O, Marzatico F, Villa RF, Dagani F, Curti D. The mitochondria electron transfer alteration as a factor involved in the brain ageing. *Neurobiol Aging*, 1992; 13: 361-8.
7. Yan LJ, Levine RL, Sohal RS. Oxidative damage during aging targets mitochondrial aconitase. *Proc Natl Acad Sci*, 1997; 94: 11168-72.
8. Benzi G, Moretti A. Age and peroxidative stress-related modifications in the cerebral enzymatic activities linked to mitochondria and glutathione system. *Free Radic Biol Med*, 1995; 19: 77-101.
9. Rego AC, Oliveira CR. Dual effect of lipid peroxidation on the membrane order of retinal cells in culture. *Arch Biochem Biophys*, 1995; 321: 127-36.
10. Makar TK, Nedergaard M, Preuss A, Gelbard AS, Perumal AS, Cooper AJL. Vitamin E, ascorbate, glutathione, glutathione disulfide, and enzymes of glutathione metabolism in culture of chick astrocytes and neurons: evidence that astrocytes play an important role in antioxidative processes in the brain. *J Neurochem*, 1994; 62: 45-53.
11. Nuttall SL, Martin U, Sinclair AJ, Kendall MJ. Glutathione in sickness and in health. *The Lancet*, 1998; 351: 645-6.
12. Pincemail J, Heusele C, Bonté F, Limet R, Defraigne JO. Stress oxydant, antioxydants nutritionnels et vieillissement. *Act Med Int*, 2001; 4, 18-23.
13. Keogh BP, Allen RG, Pignolo R, Horton J, Tresini M, Cristofalo J. Expression of hydrogen peroxide and glutathione metabolizing enzymes in human skin fibroblasts derived from donors of different ages. *J Cell Physiol*, 1996; 167: 512-22.
14. Briganti S, Cario-André M, Maaresca V, Falchi M, Urbanelli S, Taieb A, Picardo M. Relationship between skin phototype and UV-induced catalase oxidation. *Pigment Cell Res*, 2000; 13: 407.
15. Gershon D. The mitochondrial theory of ageing: Is the culprit a faulty disposal system rather than indigenous mitochondrial alterations? *Experimental Gerontology*, 1999; 34: 613-9.
16. Krutmann J. Chronically ultraviolet-exposed human skin shows a higher mutation frequency of mitochondrial DNA as compared to unexposed skin and the hematopoietic system, *Photochem Photobiol*, 1997; 66: 271-5.
17. Ray AJ, Turner R, Nikaido O, Rees JL, Birch-Machin MA. Mitochondrial DNA and Ultraviolet light exposure in human skin. *J Invest Dermatol*, 1998; 110.
18. Cardoso SM, Pereira C, Oliveira CR. Mitochondrial function is differentially affected upon oxidative stress, *Free Rad Biol Med*, 1999; 26: 3-13.
19. Gerlach M, Ben-Shachar D, Riederer P, Youdim MBH. Altered brain metabolism of iron as a cause of neurodegenerative diseases? *J Neurochem*, 1994; 63: 793-807.
20. Zorov D. Mitochondrial damage as a source of diseases and aging: a strategy of how to fight these. *Biochim. Biophys. Acta*, 1996; 1275: 10-5.
21. Kasai H, Nishimura S. Formation of 8-hydroxydeoxyguanosine in DNA by oxygen radicals and its biological significance. In: H. Sies (ed.) *Oxidative stress: oxidants and antioxidants*, Acad. Press, London, 1991; 99-116.
22. Shigenaga MK, Ames BN. Assays for 8-hydroxy-2'-deoxyguanosine, a biomarker of in vivo oxidative DNA damage. *Free Radic. Biol. Med.*, 1991; 10: 211-6.
23. Hadshiew IM, Eller MS. Age-associated decreases in human DNA repair capacity: implications for the skin. *Aging*, 1999; 22: 45-57.
24. Krutmann J, New developments in photoprotection of human skin, *Skin Pharmacol. Appl. Skin Physiol*, 2001; 14: 401-7.
25. Allen RG, Tresini M. Oxidative stress and gene regulation, *Free Rad. Biol. & Med.* 2000; 28: 463-99.
26. Dumont P, Burton M, Chen QM, Gonos ES, Fripiat C, Mazarati JB, Eliaers F, Remacle J, Toussaint O. Induction of replicative senescence biomarkers by sublethal oxidative stresses in normal human fibroblast. *Free Rad Biol & Med*, 2000; 28: 361-73.
27. Quan T, He T, Kang S, Voorhees JJ, Fisher GJ. Connective tissue growth factor: expression in human skin in vivo, and inhibition by ultraviolet irradiation. *J Invest Dermatol*, 2002; 118: 402-8.
28. Wu J, Akaike T, Hayashida K, Okamoto T, Okuyama A, Maeda H. Enhanced vascular permeability in solid tumor involving peroxynitrite and matrix metalloproteinases, *Jpn J Cancer Res*, 2001; 92: 439-51.
29. Szabo E, Virag L, BakondiE, Gyure L, Hasko G, Bai P, Hunyadi J, Gergely P, SzaboC. Peroxynitrite production,

DNA breakage, and poly(ADP-ribose) polymerase activation in a mouse model of oxazolone-induced contact hypersensitivity, *J Invest Dermatol*, 2001; 117: 74-80.

30. Augusto O, Bonini MG, Amanso AM, Linares E, Santos CC, De Menezes SL. Nitrogen dioxide and carbonate radical anion: two emerging radicals in biology, *Free Rad Biol Med*, 2002; 32: 841-59.

31. Schmid D, Muggli R, Zülfi F. Collagen glycation and skin aging. *Cosm Toil*, 2002; 118-24.

32. Rattan SIS. Synthesis, modifications and turnover of proteins during ageing. *Exp Gerontology* 1996; 31: 33-47.

33. Conconi M, Friguet B. Proteasome upon aging and on oxidation-effect of HSP 90, *Mol Biol Reports*, 1997; 24: 45-50.

34. Friguet B, Szveda LI. Inhibition of the multicatalytic proteinase (proteasome) by 4-hydroxy-2-nonenal cross-linked protein. *FEBS Letters*, 1997; 405: 21-5.

35. Bulteau AL, Moreau M, Nizard C, Friguet B. Impairment of proteasome function upon UVA- and UVB-irradiation of human keratinocytes, *Free Rad Biol Med*, 2002; 32: 1157-70.

36. Berlett BS, Stadtman ER. Protein oxidation in ageing, disease and oxidative stress. *J Biol Chem*, 1997; 272: 20313-6.

37. Stadtman ER, Berlett BS. Reactive oxygen mediated protein oxidation in aging and disease. *Chem ResTox*, 1997; 10: 485-94.

38. Dumas M, Maftah A, Bonté F, Ratinaud MH, Meybeck A, Julien R. Flow cytometric analysis of human epidermal cell ageing using two fluorescent mitochondrial probes, *C R Acad Sci Paris, Life Science*, 1995; 318: 191-7.

39. Shringarpure R, Davies KJ. Protein turnover by the proteasome in ageing and disease, *Free Rad Biol Med*, 2002; 11: 1084-9.

40. Sitte N, Merker K, Von Zglinicki T, Grune T. Protein oxidation and degradation during proliferative senescence of Human MRC-5 fibroblasts. *Free Rad Biol Med* 2000; 28 (5): 701-8.

41. Bulteau AL, Moreau M, Nizard C, Friguet B. Impairment of proteasome function upon UVA- and UVB- irradiation of human keratinocytes, *Free Rad Biol Med*, 2002; 11: 1157-70.

42. Bulteau AL, Moreau M, Nizard C, Friguet B. UVA and UVB irradiation is affecting proteasome function both in vitro and in human keratinocyte cultures. 4<sup>th</sup> Workshop on proteasome, Clermont-Ferrand, 2-4/04/2001.

43. Boudjelal M, Wang ZQ, Voorhees JJ, Fisher GJ. Ubiquitin/proteasome pathway regulates levels of retinoic acid receptor gamma and retinoid X receptor alpha in human keratinocytes. *Cancer Research*, 2000; 60: 2247-52.

44. Boudjelal M, Wang ZQ, Voorhees JJ, Fisher GJ. Ultraviolet irradiation blocks retinoid signaling in human skin through proteasome-mediated degradation of nuclear retinoid receptors. *Cancer Research*, 1998; 60: 2247-52.

45. Howard EW, Benton R, Ahern-Moore J, Tomasek J. Cellular contraction of collagen lattices is inhibited by nonenzymatic glycation. *ExpCell Res*, 1996; 228: 132-7.

46. Le Varlet B, Dumay C, Barré P, Bonté F. Influence of advanced glycation end products on the adhesion of normal human keratinocytes, *J Invest Dermatol*, 1999; 113.

47. Jeanmaire C, Danoux L, Pauly G. Glycation during human dermal intrinsic and actinic ageing: an in vivo and in vitro model study. *Brit J Dermatol*, 2001; 145: 10-8.

48. Tian WD, Gillies R, Brancalion L, Kollias N. Aging and effects of ultraviolet A exposure may be quantified by fluorescence excitation spectroscopy in vivo. *J Invest Dermatol*, 2001; 116: 840-5.

49. Vinson JA, Howard TB. Inhibition of protein glycation and advanced glycation end products by ascorbic acid and other vitamins and nutrients. *National Biochemistry*, 1996; 7: 659-63.

50. Sander CS, Chang H, Salzman S, Müller CSL, Ekanayake-Mudiyanselage S, Elsner P, Thiele JJ. Photoageing is associated with protein oxidation in human skin in vivo, *J Invest Dermatol*, 2002; 118: 618-25.

51. Scharffetter-Kochanek K, Brenneisen P, Wenk J et al. Photoageing of the skin from phenotype to mechanisms. *Exp Gerontol*, 2000; 35: 307-16.

52. Bosset S, Barré P, Bonnet-Duquennoy M, Kurfürst R, Bonté F, Schnebert S, Le Varlet B, Nicolas JF. Wrinkles: biological and immunological features, *Eur J Dermatol*, 2002; 12: 247-52.

53. Jarisch A, Krieg T, Hunzelmann N. Regulation of collagen expression by interleukin-1-β is dependent on donor age. *Acta Derm Venerol*, 1996; 76: 287-90.

54. Schmitt-Gräff A, Desmoulière A, Gabbiani G. Heterogeneity of myofibroblast phenotypic features: an example of fibroblastic cell plasticity, *Virchows Arch*, 1994; 425: 3-24.

55. Compton C, Tong Y, Trookman N, Zhao H, Roy D. TGF-β1 gene expression in cultured human keratinocytes does not decrease with biologic age. *J Invest Dermatol*, 1994; 103: 127-33.

56. Rao KMK, Cohen HJ. The role of the cytoskeleton in aging. *Exp Gerontol*, 1990; 24: 7-22.

57. Reed MJ, Ferrara NS, Vernon RB. Impaired migration, integrin function, and actin cytoskeletal organization in



- dermal fibroblasts from a subset of aged human donors. *Mechanisms of ageing and development*, 2001; 122: 1203-20.
58. Campisi J. The role of cellular senescence in skin ageing. *J Invest Dermatol Symposium Proceedings*, 1998; 3: 1-5.
59. Von Zglinicki T, Nilsson E, Döcke WD, Brunk UT. Lipofuscin accumulation and ageing of fibroblasts. *Gerontol*, 1995; 41(suppl.2): 95-108.
60. Ho HY, Cheng ML, Lu EJ, Chou YH, Stern A, Liang CM, Chiu DTY. Enhanced oxidative stress and accelerated cellular senescence in glucose-6-phosphate dehydrogenase (G6PD) deficient human fibroblasts. *Free Rad Biol Med*, 2000; 29: 156-69.
61. Reenstra WR, Yaar M, Gilchrist BA. Effect of donor age on epidermal growth factor processing in man. *Exp Cell Res*, 1993; 209: 118-122.
62. Ricciarelli R, Maroni P, Özer Nesrin, Zingg JM, Azzi A. Age-dependent increase of collagenase expression can be reduced by  $\alpha$ -tocopherol via protein kinase C inhibition. *Free Rad Biol Med*, 1999; 27: 729-37.
63. De Wit R, Capello A, Boonstra , Verkleij AJ, Post JA. Hydrogen peroxide inhibits epidermal growth factor receptor internalization in human fibroblasts. *Free Rad Biol Med*, 2000; 28: 28-38.
64. Filsell W, Rudman S, Jenkins G, Green MR. Coordinate upregulation of tenascin C expression with degree of photodamage in human skin. *Br J Dermatol*, 1999; 140: 592-9.
65. Lavrosky Y, Chatterjee B, Clark RA, Roy AK. Role of redox-regulated transcription factors in inflammation, aging and age-related diseases. *Exp Gerontol*, 2000; 35: 521-32.
66. Thibodeau A. Metalloproteinase inhibitors. *Cosmetics & Toiletries*, 2000; 115: 75-80.
67. Fenske NA, Lober CW. Continuing and functional changes of normal aging skin *J Amer Dermatol*, 1986; 15: 571-85.
68. Bologna JL. Dermatologic and cosmetic concerns the older women. *Clinic in Geriatric medicine.*, 1993; 9: 209-29.
69. Fisher GJ, Wang ZQ, Datta SC, Talwar HS, Wang ZQ, Varani J, Kang S, Voorhees JJ. Molecular basis of sun-induced premature skin ageing and retinoid antagonism. *Nature*, 1996; 379: 335-9.
70. Fisher GJ, Wang ZQ, Datta SC, Varani J, Kang S, Voorhees JJ. Pathophysiology of premature skin aging induced by ultraviolet light. *N Engl J Med*, 1997; 337: 1419-28.
71. Mauviel A. Cytokine regulation of metalloproteinase gene expression. *J Cell Biochem*, 1993; 53: 288-95.
72. Ashcroft GS, Horan MA, Herrick SE, Tarnuzzer RW, Schultz GS, Ferguson MW. Age-related differences in the temporal and spatial regulation of matrix metalloproteinases (MMPs) in normal skin and acute cutaneous wounds of healthy humans. *Cell Tissues Res*, 1997; 29, 581-91.
73. Lahmann C, Bergemann J, Harrison G, Young AR. Matrix metalloproteinase-1 and skin ageing in smokers. *The Lancet*, 2001; 357: 935-6.
74. Eisen AZ. Human skin collagenase: localization and distribution in normal human skin. *J Invest Dermatol*, 1969; 52: 442-8.
75. Bauer EA, Stricklin GP, Jeffrey JJ, Eisen AZ. Collagenase production by human skin fibroblasts. *Biochem Biophys Res Commun*, 1975; 64: 232-40.
76. Woodley DT, Kalebec T, Baner AJ, Link W, Prunieras M, Liotta L. Adult human keratinocytes migrating over nonviable dermal collagen produce collagenolytic enzymes that degrade type I and IV collagen. *J Invest Dermatol*, 1986; 86: 418-23.
77. Petersen MJ, Woodley DT, Stricklin GP, O'Keefe EJ. Constitutive production of procollagenase and tissue inhibitor of metalloproteinases by human keratinocytes in culture. *J Invest Dermatol*, 1989; 92: 156-9.
78. Petersen MJ, Woodley DT, Stricklin GP, O'Keefe EJ. Production of procollagenase by cultured human keratinocytes. *J Biol Chem*, 1987; 262: 835-40.
79. Woessner JF Jr (1991). Matrix metalloproteinases and their inhibitors in connective tissue remodeling. *FASEB J* 5: 2145-54.
80. Murphy G, Docherty AJ, Hembry RM, Reynolds JJ. Metalloproteinases and tissue damage. *Br J Rheumatol*, 1991; 30 [suppl 1]: 25-31.
81. Twining SS. Regulation of proteolytic activity in tissues. *Crit Rev Biochem Mol Biol*, 1994; 29: 315-83.
82. Fagot D, Asselineau D, Bernerd F. Direct role of human dermal fibroblasts and indirect participation of epidermal keratinocytes in MMP-1 production after UVB irradiation. *Arch Dermatol Res*, 2002; 293: 576-83.
83. Brinckmann J, Acil Y, Wolff HH, Müller PK. Collagen synthesis in (sun)-aged human skin and in fibroblasts derived from sun-exposed and sun-protected body sites. *J Photochem Photobiol*, 1995; 27: 33-8.

84. Smutzer G. Molecular demolition. *The Scientist*, 2002; 34-6.
85. Mac Cawley LJ, Matrisian LM. Matrix metalloproteinase: they're not just for matrix anymore! *Curr. Opin in Cell. Biol*, 2001, 13: 534-40.
86. Mac Cawley LJ, Matrisian LM. Matrix metalloproteinase: Multifunctional contributors to tumor progression. *Mol Med Today*, 2000; 6: 149-56.
87. Greenwald RA, Zucker S, Golub LM. Inhibition of matrix metalloproteinases: therapeutic applications. *New York Acad Sci*, 1999.
88. Bullard KM, Lund L, Mudgett JS, Mellin TN, Hunt TK, Murphy B, Werb Z, Banda MJ. Impaired wound contraction in stromelysin -1-deficient mice. *Ann Surg*, 1999; 97: 4052-7.
89. Dunsmore SE, Saarialho-Kere UK, Roby JD, Wilson CL, Matrisian LM, Welgus HG, Parks WC. Matrilysin expression and function in airway epithelium. *J Clin Invest*, 1998; 102: 1321-31.
90. Bailey AJ. Molecular mechanisms of ageing in connective tissues, *Mech. of ageing and Dev*, 2001; 122: 735-55.
91. Bornstein P. Diversity of function is inherent in matricellular proteins: an appraisal of thrombospondin 1. *J Cell Biol*, 1995; 130: 503-6.
92. Chen H, Sottile J, O'Rourke KM, Dixit VM, Mosher DF. Properties of recombinant mouse thrombospondin 2 expressed in *Spodoptera* cells. *J Biol Chem*, 1994; 271: 32226-32.
93. Chen H, Strickland DK, Mosher DF. Metabolism of thrombospondin 2. *J Biol Chem*, 1996; 271: 15993-9.
94. Yang Z, Kyriakides R, Bornstein P. Matricellular proteins as modulators of cell-matrix interactions: adhesive defect in thrombospondin 2-null fibroblasts is a consequence of increased levels of matrix metalloproteinase-2, *Mol Biol Cell*, 2000; 11: 3353-64.
95. Rieger M. Participation of Metalloproteinases in photoageing, *Allured's Cosm Toil*, 1999; 114: 65-71.
96. Xia YP, Zhao Y, Tyrone JW, Chen A, Mustoe TA. Differential activation of migration by hypoxia in keratinocytes isolated from donors of increasing age: implication for chronic wounds in the elderly. *J Invest Dermatol*, 2001; 116: 50-6.
97. Saarialho-Kere U, Kerkela E, Jeskanen L, Hasan T, Pierce R, Starcher B, Raudasoja R, Ranki A, Oikarinen A, Vaalamo M. Accumulation of matrilysin (MMP7) and macrophage metalloelastase (MMP12) in actinic damage. *J Invest Dermatol*, 1999; 113: 664-72.
98. Seo JY, Lee SH, Youn CS, Choi HR, Rhie G, Cho K, Kim KH, Park C, Eun HC, Chung JH (2001). Ultraviolet radiation increases tropoelastin mRNA expression in the epidermis of human skin in vivo. *J Invest Dermatol* 116: 915-9.
99. Tsukahara K, Takema Y, Moriwaki S, Tsuji N, Suzuki Y, Fujimura T, Imokawa G. Selective inhibition of skin fibroblast elastase elicits a concentration-dependent prevention of ultraviolet B-induced wrinkle formation. *J Invest Dermatol*, 2001; 117: 671-7.
100. Tsuji N, Moriwaki S, Suzuki S, Takema Y, Imokawa G. The role of elastases secreted by fibroblasts in wrinkle formation: implication through selective inhibition of elastase activity. *Photochem Photobiol*, 2001; 74: 283-90.
101. Watson REB, Griffiths EM, Craven NM, Shuttleworth A, Kiely CM. Fibrillin-rich microfibrils are reduced in photoaged skin. Distribution at the dermal-epidermal junction, *J Invest Dermatol*, 1999; 112: 782-7.
102. Suwabe H, Serizawa A, Kajiwara H, Ohkido M, Tsutsumi Y. Degenerative processes of elastic fibers in sun-protected and sun-exposed skin: immunoelectron microscopic observation of elastin, fibrillin-1, amyloid P, Lysozyme and  $\alpha$ 1-antitrypsin. *Pathol Internat*, 1999; 49: 391-402.
103. Gniadecka M, Nielsen OE, Wessel S, Heidenheim M, Christensen DH, Wulf HC. Water and protein structure in photoaged and chronically aged skin, *J Invest Dermatol*, 1998; 111: 1129-33.
104. Bernstein EF, Underhills CB, Hahn PJ, Brown DB, Uitto J. Chronic sun exposure alters both the content and distribution of dermal glycosaminoglycans, *J Invest Dermatol*, 1996; 135: 255-62.
105. Meyer LJM, Stern R. Age-dependent changes of hyaluronan in human skin. *J Invest Dermatol*, 1994; 102: 385-9.
106. Bonté F. Skin lipids: their origin and function. *Recent Res Devel Lipids Res*, 1999; 3: 43-62.
107. Bonté F, Saunois A, Pinguet P, Meybeck A. Existence of a lipid gradient in the upper stratum corneum and its possible biological significance. *Ach Dermatol Res*, 1997; 289: 78-82.
108. Bonté F, Pinguet P, Saunois A, Meybeck A, Beugin S, Ollivon M, Lesieur S. Thermotropic phase behavior of in vivo extracted human stratum corneum lipids, *Lip*, 1997; 32:653-60.
109. Schreiner V, Gooris GS, Pfeiffer S, Lanzendörfer G, Wenck H, Diembeck W, Proksch E, Bowstra J. Barriers characteristics of different human skin types investigated with X-ray diffraction, lipid analysis and electron microscopy imaging. *J Invest Dermatol*, 2000; 114: 654-60.
110. Sorg O, Tran C, Carraux P, Didierjean L, Falson E, Saurat JH. Oxidative stress-independent depletion of epidermal vitamin A by UVA. *J Invest Dermatol*, 2002; 118: 513-8.

111. Zouboulis CC, Boschnakov A. Chronological ageing and photoageing of the human sebaceous gland. *Clin Exp Dermatol*, 2001; 26: 600-7.
112. Dizdaroglu M, Jaruga P, Birincioglu M, Rodriguez H. Free radical-induced damage to DNA: mechanisms and measurement. *Free Rad Biol Med*, 2002; 32: 1102-15.
113. Bohr VA, Anson RM. DNA damage, mutation and fine structure DNA repair in aging. *Mutation Res*, 1995; 338: 25-34.
114. Steenken S, Jovanovic SV. How easily oxidable is DNA? One-electron reduction potentials of adenosine and guanosine radicals in aqueous solution. *J Am Chem Soc*, 1997; 119: 617-8.
115. Aruoma OI, Halliwell B. DNA and free radicals: Techniques, Mechanisms & Applications. OICA Internat. St Lucia, London, 1998; 1-20.
116. Von Zglinicki T. Stress, DNA damage and ageing – an integrative approach. *Exp Gerontol*, 2001; 36: 1049-62.
117. Cadet J, Teoule R. Comparative study of oxidation of nucleic acid components by hydroxyl radicals, singlet oxygen and superoxide anion radicals. *Photochem. Photobiol*, 1978; 28: 661-5.
118. Piette J, Calberg-Bacq CM, van de Vorst A. Alteration of guanine residues during proflavine mediated photosensitization of DNA. *Photochem. Photobiol*, 1981; 33: 325-33.
119. Kvam E, Berg K, Steen HB. Characterization of singlet-oxygen induced guanine residue damage after photochemical treatment of free nucleosides and DNA. *Biochim Biophys Acta*, 1994; 1217: 9-15.
120. Douki T, Cadet J. Modification of DNA bases by photosensitized one-electron oxidation. *Int J Radiat Biol*, 1999; 75: 571-81.
121. Ray AJ, Turner R, Nikaido O, Rees JL, Birch-Machin MA. The spectrum of mitochondrial DNA deletions is a ubiquitous marker of ultraviolet radiation exposure in human skin. *J Invest Dermatol*, 2000; 115: 674-9.
122. Pang CY, Lee HC, Yang JH, Wei YH. Human skin mitochondrial DNA deletions associated with light exposure. *Arch Biochem Biophys*, 1994; 312: 534-8.
123. Berneburg M, Gatteman N, Stege H, Grewe M, Vogelsang K, Ruzicka T, Krutmann J. Chronically UV-exposed human skin shows a higher mutation frequency of mitochondrial DNA as compared to unexposed skin and the hematopoietic system. *Photochem Photobiol*, 1997; 66: 271-5.
124. Birch-Machin MA, Tindall M, Turner R, Haldane F, Rees J. Mitochondrial DNA deletions in human skin reflect photo- rather than chronological aging. *J Invest Dermatol*, 1998; 110: 149-52.
125. Berneburg M, Grether-Beck S, Kürten V, Ruzicka T, Briviba K, Sies H, Krutmann J. Singlet oxygen mediates the UVA-induced generation of the photoageing-associated mitochondrial common deletion. *J Biol Chem*, 1999; 274 (22): 15345-49.
126. Blackburn EH. Telomere states and cell fates. *Nature*, 2000; 408.
127. Blackburn EH. Telomerase RNA structure and function. In "RNA Structure and Function" Ed. RW Simons and M Grunberg-Manago. Cold spring Harbor Laboratory Press, 1998; 669-693.
128. Blackburn EH. The telomere and telomerase: nucleic acid-protein complexes acting in a telomere homeostasis system. *Biochemistry (Moscow)*, 1997; 62: 1196-201.
129. Prescott J, Blackburn EH. Dr Jekyll or M. Hyde? Current opinion in genetics & development, 1999; 9: 368-73.
130. McEachern MJ, Krauskopf A, Blackburn EH, et al. Telomeres and their control. *Annu Rev Genet*, 2000; 34: 331-58.
131. Boukamp P. Ageing mechanisms: the role of telomere loss. *Clin Exp Dermatol*, 2001; 26: 562-5.
132. Liu JP. Studies of the molecular mechanisms in the regulation of telomerase activity. *FASEB J*. 1999; 13: 2091-104.
133. Matsui M, Miyasaka J, Hamada K, Ogawa Y, Hiramoto M, Fujimori R, Aioi A. Influence of aging and cell senescence on telomerase activity in keratinocytes. *J Dermatol Sci*, 2000; 22: 80-7.
134. Lorenz M, Saretzki, Sitte N, Metzkow S, Von Zglinicki T. BJ fibroblasts display high antioxidant capacity and slow telomere shortening independent of hTERT transfection. *Free Rad Biol Med*, 2001; 31: 824-31.
135. Yaar M, Lee MS, Rünger TM, Eller MS, Gilchrist B. Telomere mimetic oligonucleotides protect skin cells from oxidative damage. *Ann Dermatol Venerol*, 2002; 129, 1S18.
136. Kyo S, Takakura M, Kanaya T, Zhuo W, Fujimoto K, Nishio Y, Orimo A, Inoue M. Estrogen activates telomerase. *Cancer Res*, 1999; 59: 5917-21.
137. Li H, Berndt MC, Funder JW, Liu JP. Molecular interactions between telomerase and the tumor suppressor protein p53 in vitro. *Oncogene*, 1999; 18: 6785-94.
138. Ueda M, Ouhtit A, Bito T, Nakazawa K, Lübke J, Ichihashi M, Yamasaki H, Nakazawa H. Evidence for UV-associated activation of telomerase in human skin. *Cancer Res*, 1997; 57: 370-4.

139. Yin L, Hubbard AK, Giardina C. NF- $\kappa$ B regulates transcription of the mouse telomerase catalytic subunit, *J Biol Chem*, 2000; 275: 36671-5
140. Kurfürst R, Joly R, Notarnicola C, Crabbé I, André P, Bonté F, Perrier P. Epidermal prevention of premature cellular senescence by telomerase and DNA protection, 2000; *J Invest Dermatol*, 115 (3): 546
141. Farkas B, Csete B, Magyarlaki M, Bernath S, Sümegi B. Topical poly(ADP-ribose) polymerase (PARP) regulator and its prospects for use, *Ann Dermatol Venerol*, 2002; 129, 1S91.
142. Chang E, Yang J, Nagavarapu U, Herron GS. Aging and survival of cutaneous microvasculature. *J Invest Dermatol*, 2002; 118: 752-8.
143. Vijg J. Somatic mutations and aging: a re-evaluation. *Mutation Res*, 2000; 447: 117-35.
144. Haacke AR, Roublevskaia I, Cooklis M. Apoptosis: a role in skin aging? *J Invest Dermatol Symposium Proceeding*, 1998; 3: 28-35.
145. Gandarillas A. Epidermal differentiation, apoptosis and senescence: common pathway? *Exp Gerontol*, 2000; 35: 53-62.
146. Diridollou S, Vabre V, Berson M, Vaillant L, Black D, Lagarde M, Gregoire JM, Gall Y, Patat F. Skin ageing: changes of physical properties of human skin in vivo. *Internat J Cosm Sci*, 2001; 23: 353-62.
147. Tsukahara K, Takema Y, Fujimura T, Moriwaki S, Kitahara T, Imokawa G. Determination of age-related changes in the morphological structures (sagging) of the human cheek using a photometric scale and three-dimensional surface parameters. *Internat J Cosm Sci*, 2000; 22: 247-58.
148. Pellacani G, Seidenari S. Variations in facial skin thickness and echogenicity with site and age. *Acta Derm Venereol*, 1999; 79: 366-9.
149. Lévêque JL, Dresler J, Ribot-Ciscar E, Roll JP, Poelman C. Changes in tactile spatial discrimination and cutaneous coding properties by skin hydration in the elderly. *J Invest Dermatol*, 2000; 115: 454-8.
150. Baldassarri P, Calvani M. The aging process of skin and the increase in size of subcutaneous adipocytes. *Int J Tiss Reac*, 1994; 16: 229-41.
151. Quinton ND, Laird SM, Okon MA, Smith RF, Ross RJ, Blakemore AI. Serum leptin levels during the menstrual cycle of healthy women. *Br J Biomed Sci*, 1999; 56: 16-9.
152. Quinton ND, Smith RF, Clayton PE, Gills MS, Shalet S, Justice SK, Walters S, Postel-Vinay-MC, Blakemore AI, Ross RJ. Leptin binding activity changes with age: the link between leptin and puberty. *J Clin Endocrinol Metab*, 1999; 84: 2336-41.
153. Chehab FF. Leptin as a regulator of adipose mass and reproduction. *Trends in Pharmacological Sciences*, 2000; 21: 309-14.
154. Broniarczyk-Dyla G, Joss-Wichman E. Aging of the skin during menopause. *Med Sci Monit*, 1999; 5: 1024-9.
155. Bonté F. Régulations hormonales cutanées et utilisation d'extraits végétaux correcteurs par voie topique. *Phytothérapie*, 2001; 25-8.
156. Bosset S, Barré P, Chalou A, Kurfürst R, Bonté F, André P, Perrier P, Disant F, Le Varlet B, Nicolas JE. Skin ageing: clinical and histopathologic study of permanent and reducible wrinkles, *Eur J Dermatol*, 2002; 12: 247-52.
157. Talwar HS, Griffiths CEM, Fisher GJ, Hamilton TA, Voorhees JJ. Reduced type I and type III procollagens in photodamaged adult human skin. *J Invest Dermatol*, 1995; 105: 285-90.
158. Le Varlet B, Chaudagne C, Saunois, Barré P, Sauvage C, Berthoulioux B, Meybeck A, Dumas M, Bonté F. Age-related functional and structural changes in human dermo-epidermal junction components. *J Invest Dermatol Symposium Proceeding*, 1998; 3: 172-9.
159. Boorsma J, Watson REB, Craven NM, Shuttleworth CA, Kiely CM, Griffiths CEM. Collagen type VI in photoaged skin. *J Invest Dermatol*, 1996; 107.
160. Dumas M, Berville R, Barré P, Bonté, Meybeck A. In vitro processing of melanosomes by normal human keratinocytes. 18<sup>th</sup> IFSCC Congress, Venice, 1994.
161. Saunois A, Dumas M, Petit V, Bonté F. Phytochemical analysis of *Bertholletia excelsa* pericarp and its activity on fibroblasts L-ascorbic acid intracellular transport. *Rev Biol Qium*, 2001; 18 (1): 53-8.
162. Lin JY, Varani J, Kang S, Fisher GJ, Voorhees JJ. Increased stress-activated and decreased growth factor-activated MAP Kinase activities lead to collagen deficiency and reduced cell growth in skin of elderly persons. *J Invest Dermatol*, 1998; 110(4).
163. Chung JH, Kang S, Varani J, Lin J, Fischer GJ, Voorhees JJ. Decreased Extracellular-signal-regulated kinase and increased stress-activated MAP Kinase activities in aged Human skin in vivo. *J Invest Dermatol*, 2000; 114: 177-82.
164. Watson REB, Craven NM, Kang S, Jones CJP, Kiely CM, Griffiths EM. A short-term screening protocol, using fibrillin-1 as a reporter molecule, for photoaging repair agents, *J Invest Dermatol*, 2001; 116: 672-8.

165. Bykov VJ, Marcusson JA, Hemminki K. Effect of constitutional pigmentation on ultraviolet B-induced DNA damage in fair-skinned people, *J Invest Dermatol*, 2000; 114: 40-3.
166. Seiberg M. Keratinocyte-melanocyte Interactions during melanosome transfer, *Pigment Cell Res*, 2001; 14: 236-42.
167. Bandyopadhyay D, Timchenko N, Suwa T, Hornsby PJ, Campisi J, Medrano EE. The human melanocyte: a model system to study the complexity of cellular aging and transformation in non-fibroblastic cell, *Exp Ger*, 2001; 36: 1265-75.
168. Medrano E. Aging, replicative senescence, and the differentiated function of the melanocyte. The pigmenary system, Oxford University Press, 1998; 151-8.
169. Shimada M, Yamada Y, Itoh M, Yatagai T. Melanin and blood concentration in human skin studied by multiple regression analysis: experiments, *Phys Med Biol*, 2001; 46: 2385-95.
170. Shimada M, Yamada Y, Itoh M, Yatagai T. Melanin and blood concentration in human skin studied by multiple regression analysis: assessment by Monte Carlo simulation, *Phys Med Biol*, 2001; 46: 2397-406.
171. Virador VM, Muller J, Wu X, Abdel-Malek ZA, Yu ZX, Ferrans V, Kobayashi N, Wakamatsu K, Ito S, Hammer JA, Hearing VJ. Influence of alpha-melanocyte-stimulating hormone and ultraviolet radiation on the transfer of melanosomes to keratinocytes, *FASEB J*, 2002; 16: 105-7.
172. Minwalla L, Zhao Y, Le Poole C, Wickett RR, Boissy RE. Keratinocytes play a role in regulating distribution patterns of recipient melanosomes in vitro, *J Invest Dermatol*, 2001; 117: 341-7.
173. Vancoillie G, Lambert J, Mulder A, Koerten HK, Mommaas AM, Van Oostveldt P, Naeyaert JM. Kinesin and kinectin can associate with the melanosomal surface and form a link with microtubules in normal human melanocytes, *J Invest Dermatol*, 2000; 114: 421-9.
174. Hara M, Yaar M, Byers R, Goukassian D, Fine RE, Gonsalves J, Gilchrist BA. Kinesin participates in melanosomal movement along melanocyte dendrites, *J Invest Dermatol*, 2000; 114: 438-43.
175. Scott G, Zhao Q. Rab3 and SNARE proteins: potential regulators of melanosome movement. *J Invest Dermatol*, 2001; 116: 296-304.
176. Seiberg M, Paine C, Sharlow E, Andrade-Gordon P, Costanzo M, Eisinger M, Shapiro SS. Inhibition of melanosome transfer results in skin lightening. *J Invest Dermatol*, 2000; 115: 162-7.
177. Manaka I, Kadono S, Kawashima M, Kobayashi T, Imokawa G. The mechanisms of hyperpigmentation in seborrheic keratosis involves the high expression of endothelin-converting enzyme-1 $\alpha$  and TNF- $\alpha$ , which stimulate secretion of endothelin 1, *Brit J Dermatol*, 2001; 145: 895-903.
178. Tajima S, Manaka I, Kawashima M, Kobayashi T, Imokawa G. Role of endothelin cascade between keratinocytes and melanocytes in hyperpigmentation in senile freckles, *J Invest Dermatol*, 1998; 110 (4).
179. Craven NM, Helbling I, Chadwick CA, Ferguson JE, Potten CS, CEM Griffiths. Photoageing is associated with persistence of thymine dimers in epidermis and dermis. *Brit J Dermatol*, 1998; 138: 724-53.
180. Nofsinger JB, Liu Y, Simon JD. Aggregation of eumelanin mitigates photogeneration of reactive oxygen species. *Free Rad Biol Med*, 2002; 32: 720-30.
181. Andersen WK, Labadie RR, BS, Bhawan J, MD. Histopathology of solar lentiginosities of the face: a quantitative study. *J Am Acad Dermatol*, 1997; 36: 444-7.
182. Fushimi H, Shishido E, Kawashima M, Ichikawa M, Imokawa G. The role of the stem cell factor (CF) in hyperpigmentation mechanism involved in senile freckles (SF), *Pigment Cell Res*, 2000; 13: 415.
183. Higuchi K, Kawashima M, Ichikawa Y, Imokawa G. Effects of sphingosylphosphorylcholine (SPC) on proliferation and melanogenesis in human melanocytes, *Pigment Cell Res*, 2000; 13: 415.
184. Hoogduijn M, Ancans A, Thody AJ. Mammalian pigment cells express the MCH receptor, *Pigment Cell Res*, 2000; 13: 400.
185. Thody AJ. Hormonal control of skin pigmentation, *Pigment Cell Res*, 2000; 13: 404.
186. Sasaki M, Horikoshi T, Kizawa K, Igarashi S, Uchiwa H, Miyachi Y. Suppression of melanogenesis by induction of metallothionein in human melanocytes, *Pigment Cell Res*, 2000; 13: 411.
187. Evans SK, Lundblad V. Positive and negative regulation of telomerase access to the telomere. *J Cell Sci*, 2000; 113: 3357-64.
188. Colgin L, Wilkinson C, Engelow A, et al. The hTERT alpha splice variant is a dominant negative inhibitor of telomerase activity. *Neoplasia*, 2000; 2: 426-32.
189. Stampfer MR, Garbe J, Levine G, et al. Expression of the telomerase catalytic subunit, hTERT, induces resistance to transforming growth factor beta growth inhibition in p16INK4A (-) human mammary epithelial cells. *Proc Natl Acad Sci. USA*, 2001; 98: 4498-503.

190. Beattie TL, Zhou W, Robinson MO, et al. Functionnal multimerization of the human telomerase reverse transcriptase, *Mol Cemll Biol*, 2001; 21: 151-60.
191. Hillebrand GG, Levine MJ, Miyamoto K., The age-dependent changes in skin conditions in African Americans, Asian Indians, Caucasians, East Asians and Latinos. *IFSCC Magazine*, 2001; 4: 259-66.
192. Nagashima H, Hanada K, Hashimoto I. Correlation of skin phototype with facial wrinkle formation. *Photodermatol Photoimmunol Photomed*, 1999; 15: 2-6.
193. Camel E, Arnaud-Boissel L, Schnebert S, Neveu M, Tan SK, Guillot JP. Does Asian skin induce significant changes in sun protection factor (SPF) determination compared to Caucasian skin: one of the first in vivo correlations, *IFSCC Magazine*, 2002; 5: 31.
194. Wee LKS, Chong TK, Koh Soo Quee D. Assessment of skin types, skin colours and cutaneous responses to ultraviolet radiation in an Asian population, *Photodermatol Photoimmunol Photomed*, 1997; 13: 169-72.
195. Matsumoto M, Kobayashi N, Hoshina O, Hayashi S, Arai S, Study of causal factors of dark circles around th eyes, *IFSCC Magazine*, 2001; 4: 281.
196. Le Fur I, Numagami K, Guinot C, Lopez S, Morizot F, Lambert V, Kobayashi H, Tschachler E, Tagami H, Age-related reference values of skin colour in Caucasian and Japanese healthy women according to skin site, XXIth IFSCC International Congress 2000, Berlin.
197. Sakamaki T, Hayashi S, Hayakawa R. Melanin distribution in corneocytes and fine patterns of pigmentation on pigmented areas in pigmentary disorders, *Skin Res Technol*, 1999; 5: 37-41.
198. Tsukahara K, Takema Y, Fujimura T, Moriwaki S, Kitahara T, Imokawa G. Age-related alterations of echogenicity in Japanese skin, *Dermatol*, 2000; 200: 303-7.
199. Park SG, Kim YD, Kim JJ, Kang SH, Two possible classifications of facial skin type by two parameters in Korean women: sebum excretion rate (SER) and skin surface relief (SSR), *Skin Res Technol*, 1999; 5: 189-94.
200. Yamazaki S, Ozawa N, Hiratsuka A, Watabe T. Increases in cholesterol 7-hydroxyperoxides in lipids of human skin by sunlight exposure, *Free Rad Biol Med*, 1999; 26: 1126-33
201. Chung JH, Lee SH, Youn CS, Park BJ, Kim KH, Park KC, Cho KB, Eun HC. Cutaneous photodamage in Koreans: influence of sex, sun exposure, smoking, and skin color, *Arch Dermatol*, 2001; 137: 1043-51.
202. Mangubat MI, Estanislao R, Suero M, Rivera Z, Li J, Watanabe H, Hyun OK. Characterization of Asian Skin (Part): Influence of age, *Ann Dermatol Venerol*, 2002; 129: 1S402.
203. Won CH, Youn CS, Chung JH, Kim KH, Cho KH, Eun HC. The endocrinological effects on wrinkle in Korean Women, *Ann Dermatol Venerol*, 2002; 129: 1S406.
204. Yang ZL, D Lacharrière O, Li L, Lian S, Yang FZ, Zou W, Nouveau S, Qian BQ, Bouillon C, Ran YP. Facial wrinkles in Chinese skin: impact of climatic factors. A clinical study on 2, 000 Chinese women, *Ann Dermatol Venerol*, 2002; 129: 1S112.
205. Yang ZL, D Lacharrière O, Nouveau-Richard S, Mac Mary S, Bastien P, Humbert P. Skin aging. A comparison between Chinese and European population. *Ann Dermatol Venerol*, 2002; 129: 1S112.
206. Li L, De Lacharrière O, Lian S, Yang ZL, Yang FZ, Zhou W, Nouveau-Richard S, Qian BY, Bouillon C, Ran YP. Pigmented spots on face and hands: specific features in Chinese skin. A clinical study on 2, 000 Chinese women, *Ann Dermatol Venerol*, 2002; 129: 1S111.

**A U T H O R S** ' Céline Rocquet, LVMH R&D - Parfums et Cosmétiques, 185 Avenue de  
**A D D R E S S E S** Verdun, 45804 Saint Jean de Braye, France  
Frédéric Bonté, PhD, biochemist, same address, fbonte@diormail.com